

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 369 431 A1**

(12)

**EUROPEAN PATENT APPLICATION**

published in accordance with Art. 158(3) EPC

(43) Date of publication:

**10.12.2003 Bulletin 2003/50**

(21) Application number: **02801565.9**

(22) Date of filing: **15.10.2002**

(51) Int Cl.7: **C07K 16/28**, C12N 5/10,  
C12N 15/13, C12N 1/21,  
C12N 1/19, C12P 21/08,  
A61K 39/395

(86) International application number:  
**PCT/JP02/10665**

(87) International publication number:  
**WO 03/033538 (24.04.2003 Gazette 2003/17)**

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
IE IT LI LU MC NL PT**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **15.10.2001 JP 2001317054**

(71) Applicant: **KIRIN BEER KABUSHIKI KAISHA  
Tokyo 104-8288 (JP)**

(72) Inventors:

- **TAWARA, Tomonori**,  
c/o Pharmaceutical Research Lab  
Takasaki-shi, Gunma 370-1295 (JP)
- **KATAOKA, Shiro**,  
c/o Pharmaceutical Research Labor.  
Takasaki-shi, Gunma 370-1295 (JP)

(74) Representative: **HOFFMANN - EITLE  
Patent- und Rechtsanwälte  
Arabellastrasse 4  
81925 München (DE)**

(54) **ANTI-HLA-DR ANTIBODY**

(57) This invention provides an anti-HLA-DR monoclonal antibody. This invention relates to an antibody binding to HLA-DR or a functional fragment thereof having (a) life-extending effects in nonhuman animals bearing HLA-DR-expressing cancer cells and (b) activity of suppressing immune responses lower than that of L243,

or an antibody binding to HLA-DR or a functional fragment thereof exhibiting immunosuppressive activity equivalent to or higher than that of the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55).

**EP 1 369 431 A1**

**Description**

## Technical Field

5 **[0001]** The present invention relates to an anti-HLA-DR antibody that recognizes the human leucocyte antigen-DR (HLA-DR), which is a cell membrane molecule associated with immunity. Further, the present invention relates to the following two types of agents comprising, as an active ingredient, the anti-HLA-DR antibody: (1) a preventive or therapeutic agent for diseases caused by an HLA-DR-expressing cell, especially, a therapeutic agent for malignant tumors; and (2) a preventive or therapeutic agent for immune responses caused by an HLA-DR-expressing cell, especially, a therapeutic agent for chronic rheumatism.

## Background Art

15 **[0002]** The use of an antibody, which binds to a protein expressed on a cell surface and is capable of leading the cell to death or toxicity, has been attempted in the treatment of cancer, etc. At present, a chimeric antibody (Rituximab) targeting CD20, which is a receptor existing on a cell membrane, and a monoclonal antibody such as a humanized antibody targeting Her2/neu are used in the treatment of malignant tumors, and their therapeutic effects are acknowledged. An antibody is characterized by a long serum half-life and high specificity for an antigen, and thus is particularly useful as an anti-tumor agent. For example, when an antibody targeting a tumor-specific antigen is administered, accumulation thereof in the tumor is presumed. Thus, an attack by the immune system due to the complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC) on cancer cells can be expected. Binding of a radionuclide or an agent such as a cytotoxic substance to the antibody enables the effective transmission of the bound agent to a tumor site. This also reduces the amount of the agent reaching other non-specific tissues, and thus reduced side effects can be expected. When a tumor-specific antigen has activity for inducing cell death, an agonistic antibody is administered. In contrast, when a tumor-specific antigen is associated with growth and survival of cells, a neutralizing antibody is administered. This can result in the accumulation of tumor-specific antibodies and arrest or regression of tumor growth due to the activity of the antibody. As mentioned above, antibodies are considered suitable for application as anti-tumor agents because of their features.

20 **[0003]** Recently, significant anti-tumor effects of Rituximab have been exhibited with respect to B-cell lymphoma, and the side effects thereof are limited. Since Rituximab is a chimeric human-mouse protein, however, the antigenicity of Rituximab itself is strong, an antibody against a mouse moiety is produced inside the body, and the effect could be deteriorated. For some types of cancer, the therapeutic effect of Rituximab is low with the use of Rituximab alone, and the combined use thereof with an anticancer agent is currently being clinically examined (see McLaughlin P. et. al., J Clin Oncol. (1998), 16, 2825-2833; Coiffier B. et al., Blood (1998), 92, 1927-81). Accordingly, a novel anti-tumor antibody targeting an antigen is needed, and a monoclonal antibody against HLA-DR, which is a class II major histocompatibility complex (MHC) molecule, can be expected to have clinical anti-tumor activity as an antibody that recognizes an antigen different from that recognized by Rituximab.

25 **[0004]** In contrast, class II major histocompatibility complex (MHC) molecules bind to antigen peptide fragments and present these antigen peptide fragments to helper (CD4<sup>+</sup>) T-cells ("Th" cells) (see Babbin B. et al., Nature (1985), 317, 359-361). A monoclonal antibody that is specific for the class II MHC molecules is reported as a very potent selective inhibitor against the immune response of Th cells *in vitro* (see Baxevanis CN, et. al., Immunogenetics (1980), 11, 617-625). Since this monoclonal antibody was discovered, it has been considered to be an agent that can be used in the selective immunosuppressive therapy of autoimmune diseases such as chronic rheumatism. Based on the initial *in vivo* research, significant effects of these monoclonal antibodies on Th cellular heterogeneity and autoimmune response have been elucidated (see Rosenbaum JT. et al., J. Exp. Med. (1981), 154, 1694-1702; Waldor MK. et al., Proc. Natl. Acad. Sci. USA (1983), 80, 2713-2717; Jonker M. et al., J. Autoimmun. (1988), 1, 399-414; Stevens HP. et al., Transplant. Proc. (1990), 22, 1783-1784). Further, as a result of research using primates, it was discovered that graft-versus-host disease in homograft was suppressed (Billing R. & Chatterjee S. (1983), Transplant. Proc., 15, 649-650; Jonker M. et al., Transplant Proc. (1991), 23, 264-265).

30 **[0005]** Currently, immunological rejection at the time of organ transplantation is clinically suppressed using immunosuppressive agents such as cyclosporin A or FK506. A disadvantage of these immunosuppressive agents is that potent side effects are caused by the non-specific suppression of the immune response.

35 **[0006]** Accordingly, antibodies are considered suitable for use as immunosuppressive agents with few side effect because of their features.

## Disclosure of the Invention

40 **[0007]** An immunosuppressive agent using a human antibody having high immunosuppressive activity and low im-

munogenicity has not yet been developed.

**[0008]** An object of the present invention is to produce such an antibody and use it as an anti-tumor agent or immunosuppressive agent

**[0009]** The present inventors have conducted concentrated studies in order to produce an antibody against human HLA-DR. As a result, they have succeeded in obtaining a monoclonal antibody exhibiting an anti-tumor effect at very low concentration on HLA-DR-expressing cancer cells and a monoclonal antibody that specifically suppresses immune activity through HLA-DR. Further, they have identified the sequence in the variable region of the monoclonal antibody and determined the epitope to which the monoclonal antibody binds. This has led to the completion of the present invention.

**[0010]** More specifically, the present invention is as follows.

**[0011]** The present invention provides, in the first aspect thereof, a monoclonal antibody that binds to HLA-DR produced from a mouse-mouse hybridoma, for example, a monoclonal antibody that is preferably a human antibody produced from HD4, HD6, HD8, or HD10, or a functional fragment thereof. The monoclonal antibody produced from HD4, HD6, HD8, or HD10 is of an immunoglobulin G (IgG) type. The hybridoma HD8 and the hybridoma HD10 are deposited internationally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of October 11, 2001 under the accession numbers FERM BP-7773 and FERM BP-7774, respectively. The hybridoma HD4 is deposited internationally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of October 11, 2001 under the accession number FERM BP-7771. The hybridoma HD6 is deposited internationally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of October 11, 2001 under the accession number FERM BP-7772.

**[0012]** According to an embodiment of the present invention, the antibody according to the present invention comprises a variable region of the antibody produced from the aforementioned hybridoma or a functional fragment thereof.

**[0013]** In another embodiment of the present invention, the antibody according to the present invention includes an antibody having a modified subclass, which is an antibody produced from the hybridoma HD8 having an IgG1, IgG2, IgG3, or IgG4 subclass or a functional fragment thereof, an antibody produced from the hybridoma HD4 having an IgG1, IgG2, IgG3, or IgG4 subclass or a functional fragment thereof, an antibody produced from the hybridoma HD10 having an IgG1, IgG2, IgG3, or IgG4 subclass or a functional fragment thereof, or an antibody produced from the hybridoma HD6 having an IgG1, IgG2, IgG3, or IgG4 subclass or a functional fragment thereof. According to a further embodiment of the present invention, the antibody according to the present invention is an antibody having a modified amino acid sequence in the constant region of the heavy chain or a functional fragment thereof. For example, an antibody comprises amino acid 331 in the constant region of the heavy chain according to the EU numbering system (see Sequences of Proteins of Immunological Interest, NIH Publication No. 91-3242) being substituted with Ser or a functional fragment thereof.

**[0014]** In another embodiment of the present invention, the subclass of the antibody or a functional fragment thereof is rearranged to IgG1, IgG2, or IgG4, amino acid 331 in the constant region of the heavy chain according to the EU numbering system is substituted with Ser, and the subclass is modified as IgG1, IgG1Ser, IgG2, IgG2Ser, or IgG4. As a result, only an antibody or a functional fragment thereof having an IgG1 or IgG1Ser subclass develops ADCC, and only an antibody or a functional fragment thereof having an IgG1 or IgG2 subclass develops CDC activity.

**[0015]** In another aspect of the present invention, the present invention provides an antibody that binds to HLA-DR or a functional fragment thereof comprising a variable region of an antibody produced from the hybridoma HD4, HD6, HD8, or HD10. In an embodiment of the present invention, the antibody or a functional fragment thereof according to the present invention comprises a variable region of an antibody produced from the hybridoma HD8 having an amino acid sequence in the mature variable region of the amino acid sequences as shown in SEQ ID NOs: 21 and 23. In another embodiment of the present invention, the antibody or a functional fragment thereof according to the present invention comprises a variable region of an antibody produced from the hybridoma HD4 having the amino acid sequence in the mature variable regions in the amino acid sequences as shown in SEQ ID NOs: 17 and 19.

**[0016]** In a further aspect of the present invention, the present invention relates to an antibody or a functional fragment thereof that can bind to a specific epitope of HLA-DR. In embodiments of the present invention, the antibody according to the present invention is an antibody that binds to HLA-DR or a functional fragment thereof, which maximally binds to the peptide as shown in SEQ ID NO: 82. This peptide is selected from among peptides that are prepared by shifting 2 amino acids of the amino acids in the extracellular region (the amino acid sequence being shown in SEQ ID NO: 147 and the nucleotide sequence being shown in SEQ ID NO: 146) of the HLA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides (preparing 13-mer peptides for 199 amino acids, i.e., amino acids 29 to 227, in the amino acid sequence as shown in SEQ ID NO: 147), binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus. Also, the antibody according to the present invention is an antibody that binds to HLA-DR or a functional fragment thereof, which potently binds to all three peptides as shown in SEQ ID NOs: 82, 83, and 84.

These peptides are selected from among peptides that are prepared by shifting 2 amino acids of the amino acids in the extracellular region of the HLA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides, binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus. Further, the antibody according to the present invention is an antibody that binds to HLA-DR or a functional fragment thereof, which significantly binds to all the peptides as shown in SEQ ID NOs: 24 to 39 and all the peptides as shown in SEQ ID NOs: 40 to 43. These peptides are prepared by shifting 2 amino acids of the amino acids in the extracellular region of the "LA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides, binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus.

**[0017]** Furthermore, the present invention provides, in another aspect, an antibody or a functional fragment thereof that can extend the survival of individual mice by suppressing tumor growth (for example, those derived from the Raji cell transplanted to SCID mice). The amount of the antibody or a functional fragment thereof according to the present invention to be administered to tumor-bearing test animals (for example, tumor-bearing test animals such as lymphoma cell-bearing mouse models, each with a body weight of 20 g) is 0.1  $\mu\text{g}/\text{body}$  to 1  $\mu\text{g}/\text{body}$  or 5  $\mu\text{g}/\text{kg}$  to 50  $\mu\text{g}/\text{kg}$ . Examples of dose are 1  $\mu\text{g}/\text{body}$  or 50  $\mu\text{g}/\text{kg}$ , and preferably 0.1  $\mu\text{g}/\text{body}$  or 5  $\mu\text{g}/\text{kg}$ .

**[0018]** In an embodiment of the present invention, the antibody according to the present invention binds to HLA-DR or a functional fragment thereof, which has the following properties (a) and (b):

(a) when a 6-week-old SCID mouse is inoculated intravenously with 10  $\mu\text{L}$  of the anti-asialo GM1 antiserum, on the next day, inoculated intravenously with  $5 \times 10^6$  of Burkitt's lymphoma cells Raji (ATCC CCL-86), and 5 days thereafter, inoculated with 5 to 50  $\mu\text{g}/\text{kg}$ , based on body weight, of the antibody of the present invention, the survival ratio of the mouse is higher than that achieved when inoculated with the same amount of the human anti-HSA antibody; and

(b) the immunosuppressive activity is lower than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) at the same concentration, wherein the immunosuppressive activity is assayed as follows: 50  $\mu\text{L}$  of antibody adjusted at 8  $\text{mg}/\text{mL}$ , and preferably 8  $\text{mg}/\text{mL}$  using 10% FCS-containing RPMI 1640 medium is mixed with 50  $\mu\text{L}$  of mature dendritic cell suspension derived from a first human donor adjusted at  $2 \times 10^5$  cells/mL using 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate, the mixture is allowed to stand at 4°C for 30 minutes, the resultant is mixed with 100  $\mu\text{L}$  of T-cell suspension (purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using 10% FCS-containing RPMI 1640 medium derived from a second human donor having a histocompatible antigen different from that of the first human donor, the mixture is cultured at 37°C in the presence of 5%  $\text{CO}_2$  for 5 days,  $^3\text{H}$  thymidine is added thereto at 1.0  $\mu\text{Ci}/\text{well}$ , the resultant is cultured at 37°C in the presence of 5%  $\text{CO}_2$  for 16 to 20 hours, the  $^3\text{H}$  thymidine incorporated in the cell is recovered and then measured using a scintillator, and the incorporation of the  $^3\text{H}$  thymidine into the cell is used as an indicator to assay the immunosuppressive activity.

**[0019]** Further, the present invention provides, in another aspect thereof, an antibody recognizing HLA-DR or a functional fragment thereof, which exhibits immunosuppressive activity equivalent to or higher than that of the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55). In an embodiment of the present invention, the antibody or a functional fragment thereof according to the present invention has the immunosuppressive activity equivalent to or higher than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) at the same concentration, wherein the immunosuppressive activity is assayed in the following manner. First, 50  $\mu\text{L}$  of antibody adjusted at 8  $\text{mg}/\text{mL}$  using 10% FCS-containing RPMI 1640 medium is mixed with 50  $\mu\text{L}$  of mature dendritic cell suspension derived from a first human donor adjusted at  $2 \times 10^5$  cells/mL using 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate. The mixture is then allowed to stand at 4°C for 30 minutes, and the resultant is mixed with 100  $\mu\text{L}$  of T-cell suspension (purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using 10% FCS-containing RPMI 1640 medium derived from a second human donor having a histocompatible antigen different from that of the first human donor. The resultant is cultured at 37°C in the presence of 5%  $\text{CO}_2$  for 5 days,  $^3\text{H}$  thymidine is added thereto at 1.0  $\mu\text{Ci}/\text{well}$ , the resultant is further cultured at 37°C in the presence of 5%  $\text{CO}_2$  for 16 to 20 hours, the  $^3\text{H}$  thymidine incorporated in the cell is recovered and then measured using a scintillator, and incorporation of the  $^3\text{H}$  thymidine into the cell is used as an indicator to assay the immunosuppressive activity.

**[0020]** The present invention further provides, in another aspect, a nucleic acid encoding an antibody comprising a variable region of an antibody produced from a hybridoma or a functional fragment thereof, wherein said nucleic acid is possessed by a hybridoma selected from the group consisting of the hybridoma HD8 (accession number FERM BP-7773), the hybridoma HD10 (accession number FERM BP-7774), the hybridoma HD4 (accession number FERM BP-7771), and the hybridoma HD6 (accession number FERM BP-7772), a protein encoded by the nucleic acid, an expression vector having the nucleic acid, and a host selected from the group consisting of *E. coli*, yeast cell, insect cell, mammalian cell, plant cell, and mammals having the expression vector. The present invention provides in its embodiments: a nucleic acid encoding the antibody or a functional fragment thereof, which comprises a variable region

having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 17 and 19; a nucleic acid encoding the antibody or a functional fragment thereof, which comprises a variable region having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 21 and 23; and a nucleic acid encoding the antibody or a functional fragment thereof, wherein the antibody is selected from the group consisting of the antibody HD8G1Ser, the antibody HD8G2Ser, and the antibody HD4G2Ser. The antibody HD8G1Ser is the antibody HD8 having an IgG1 subclass and amino acid 331 according to the EU numbering system being substituted with Ser, the antibody HD8G2Ser is the antibody HD8 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser, and the antibody HD4G2Ser is the antibody HD4 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser.

**[0021]** The present invention further provides, in another aspect thereof, a process for producing the anti-HLA-DR monoclonal antibody, wherein a gene encoding the anti-HLA-DR monoclonal antibody is isolated from a hybridoma selected from the group consisting of the hybridoma HD8 (accession number FERM BP-7773), the hybridoma HD10 (accession number FERM BP-7774), the hybridoma HD4 (accession number FERM BP-7771), and the hybridoma HD6 (accession number FERM BP-7772), an expression vector having said gene is constructed, the expression vector is introduced into a host to express the monoclonal antibody, and the anti-HLA-DR monoclonal antibody is collected from the resulting host, a culture supernatant of the host, or a secretion product of the host.

**[0022]** The present invention further provides, in another aspect thereof, a preventive, therapeutic, or diagnostic agent for tumors, which comprises, as an active ingredient, the aforementioned antibody or a functional fragment thereof.

**[0023]** Examples of tumors that can be prevented or treated include at least one member selected from the group consisting of leukemia (including chronic lymphatic leukemia and acute lymphatic leukemia), lymphoma (including non-Hodgkin's lymphoma, Hodgkin's lymphoma, T-cell lymphoma, B-cell lymphoma, Burkitt's lymphoma, malignant lymphoma, diffuse lymphoma, and follicular lymphoma), myeloma (including multiple myeloma), breast cancer, colon cancer, kidney cancer, gastric cancer, ovarian cancer, pancreatic cancer, cervical cancer, endometrial cancer, esophageal cancer, liver cancer, head and neck squamous cancer, skin cancer, urinary tract cancer, prostate cancer, choriocarcinoma, pharyngeal cancer, laryngeal cancer, pleural tumor, arrhenoblastoma, endometrial hyperplasia, endometriosis, embryoma, fibrosarcoma, Kaposi's sarcoma, angioma, cavernous angioma, hemangioblastoma, retinoblastoma, spongiocytoma, neurofibroma, oligodendroglioma, medulloblastoma, neuroblastoma, neuroglioma, rhabdomyoblastoma, glioblastoma, osteogenic sarcoma, leiomyosarcoma, thyroid sarcoma, and Wilms tumor.

**[0024]** The present invention provides, in another aspect, an immunosuppressive agent comprising, as an active ingredient, the antibody or a functional fragment thereof according to the present invention. The present invention further provides a preventive, therapeutic, or diagnostic agent for autoimmune diseases or allergies, which comprises, as an active ingredient, the antibody or a functional fragment thereof according to the present invention.

**[0025]** In an embodiment of the present invention, a preventive or therapeutic agent is an immunosuppressive agent at the time of organ transplantation (a preventive or therapeutic agent for immunological rejection at the time of pancreatic islet or kidney transplantation or GVHD), a therapeutic agent for autoimmune diseases (for example, rheumatism, arteriosclerosis, multiple sclerosis, systemic erythematodes, idiopathic thrombocytopenia, or Crohn's disease), or a preventive or therapeutic agent for allergic diseases such as asthma. In another embodiment of the present invention, the life-extending effects are recognized in the tumor-bearing SCID mice to which the Raji cell had been transplanted 5 days after the tumor transplantation with the administration of 5 µg/kg or lower of the antibody or a functional fragment thereof according to the present invention.

**[0026]** The present invention includes an antibody or a functional fragment comprising the amino acid sequences in the mature variable region of the heavy chain and that of the light chain of the antibody produced from the hybridoma HD4 as shown in SEQ ID NO: 17 or 19 and the amino acid sequences in the mature variable region of the heavy chain and that of the light chain of the antibody produced from the hybridoma HD8 as shown in SEQ ID NO: 21 or 23.

**[0027]** The aforementioned antibody or a functional fragment thereof comprises, for example, the amino acid sequence in the mature variable region of the heavy chain and that of the light chain encoded by the nucleic acid sequence isolated from the hybridoma HD4 as shown in SEQ ID NO: 16 or 18 and the amino acid sequence in the mature variable region of the heavy chain and that of the light chain encoded by the nucleic acid sequence isolated from the hybridoma HD8 as shown in SEQ ID NO: 20 or 22.

**[0028]** The present invention is hereafter described in detail.

**[0029]** It is also reported that the anti-HLA-DR monoclonal antibody has activity of suppressing immune responses. Based on the initial *in vivo* research, significant effects of the anti-HLA-DR monoclonal antibody on Th cellular heterogeneity and autoimmune response were elucidated (see Rosenbaum JT. et al., J. Exp. Med. (1981), 154, 1694-1702; Waldor MK. et al., Proc. Natl. Acad. Sci. USA (1983), 80, 2713-2717; Jonker M. et al., J. Autoimmun. (1988), 1, 399-414; Stevens HP. et al., Transplant. Proc. (1990), 22, 1783-1784). Further, as a result of research using primates, it was discovered that graft-versus-host disease in homograft was suppressed (Billing R. & Chatterjee S. (1983), Transplant. Proc., 15, 649-650; Jonker M. et al., Transplant Proc. (1991), 23, 264-265). These reported antibodies, however, are

mouse antibodies. Recently, Protein Design Labs Inc. has developed a humanized HLA-DR antibody using the mouse anti-HLA-DR antibody 1D10 and converting regions other than the variable region into the sequence of the human antibody by their humanizing techniques and gene recombination (see Sheri AK. et. al., Int. J. Cancer (2001), 93, 556-565). This is clinically examined in the U.S.

**[0030]** The novel human anti-HLA-DR monoclonal antibody according to the present invention is a complete human antibody, and the antigenicity against the mouse sequences, which is always problematic in the mouse antibody, has already been resolved.

**[0031]** Any of the immunoglobulin G (IgG), immunoglobulin A (IgA), immunoglobulin E (IgE), and immunoglobulin M (IgM) antibodies can be suitably used. In general, IgG is preferable.

**[0032]** The terms used in the present invention are defined in order to describe the present invention in more detail.

#### 1. HLA-DR and an antibody thereof

**[0033]** The antibody according to the present invention is an antibody against HLA-DR, which is a class II major histocompatibility complex (MHC), i.e., an antibody that recognizes and binds to HLA-DR and an antibody that has reactivity with HLA-DR.

**[0034]** The "antibody that binds to HLA-DR" in the present invention is an antibody or a portion thereof having reactivity with human HLA-DR or a portion thereof, and it includes a functional fragment thereof. A "functional fragment" refers to a portion of an antibody (a partial fragment), which has at least one function of the antibody on an antigen. Specific examples thereof include F(ab')<sub>2</sub>, Fab', Fab, Fv, disulphide-linked Fv, single-chain Fv (scFv), and a polymer thereof (D. J. King, Applications and Engineering of Monoclonal Antibodies, 1998, T. J. International Ltd.). Or, a "functional fragment" is a fragment of an antibody that is capable of binding to an antigen. A functional fragment of the antibody according to the present invention binds to HLA-DR and exhibits an anti-tumor effect or potent immunosuppressive activity.

**[0035]** The term "human antibody" used herein refers to an antibody, which is an expression product of a human-derived antibody gene. The human antibody can be obtained by introducing a human antibody locus and administering an antigen to a transgenic animal that is capable of producing a human-derived antibody as described below. An example of such a transgenic animal is a mouse, and a process for producing a mouse that can produce a human antibody is described in WO 02/43478.

**[0036]** Examples of the antibody according to the present invention include various antibodies exhibiting anti-tumor effects at low concentration on human HLA-DR-expressing cancer cells as described in the examples below.

**[0037]** The antibody according to the present invention includes a monoclonal antibody comprising the heavy chain and/or light chain having an amino acid sequence with deletion, substitution, or addition of one or several amino acids in various amino acid sequences thereof. The aforementioned partial modification of amino acid (deletion, substitution, insertion, or addition) can be introduced into the amino acid sequence of the antibody according to the present invention by partially modifying the nucleotide sequence encoding the amino acid sequence of interest. Such partial modification of the nucleotide sequence can be introduced by a general method of conventional site specific mutagenesis (Proc Natl Acad Sci USA, 1984, Vol. 81: 5662). The antibody used herein refers to an immunoglobulin in which all regions including a variable region and a constant region of the heavy chain and a variable region and a constant region of the light chain constituting the immunoglobulin are derived from a gene encoding the immunoglobulin.

**[0038]** The antibody according to the present invention includes an antibody having any immunoglobulin class and isotype.

**[0039]** The anti-HLA-DR antibody according to the present invention can be produced by a process as described below. Specifically, human HLA-DR, a part thereof, a binding product thereof with a suitable carrier substance for enhancing antigenicity of the antigen (e.g., bovine serum albumin), or the like is administered to a non-human mammalian, such as a human antibody-producing transgenic mouse, for immunization together with an immunopotentiating agent (e.g., Freund's complete or incomplete adjuvant), if necessary. Alternatively, a gene encoding the human HLA-DR  $\alpha$  chain or  $\beta$  chain can be introduced, and an animal cell having excessively expressed HLA-DR on its surface can be administered for immunization. A monoclonal antibody can be obtained by culturing a hybridoma obtained by fusing an antibody-producing cell obtained from the immunized animal and a myeloma cell incapable of producing an autoantibody, and selecting a clone that produces a monoclonal antibody having specific affinity to the antigen used for the immunization.

**[0040]** The antibody according to the present invention includes those having different subclasses that were modified by genetic engineering known to a person skilled in the art (see, for example, EP 314161). Specifically, an antibody having a subclass that is different from the original subclass can be obtained using DNA encoding a variable region of the antibody according to the present invention using a genetic engineering technique. For example, the subclass of the antibody according to the present invention can be converted into IgG2 or IgG4 to obtain an antibody having a low degree of binding to the Fc receptor. On the contrary, the subclass of the antibody according to the present invention

can be converted into IgG1 or IgG3 to obtain an antibody having a high degree of binding to the Fc receptor. Further, modification of the amino acid sequence in the constant region of the antibody according to the present invention by genetic engineering or binding with a sequence of a constant region having such a sequence enables changing of the degree of binding to the Fc receptor (see Janeway CA. Jr. and Travers P. (1997), Immunobiology, Third Edition, Current Biology Ltd./Garland Publishing Inc.) or that to the complement (see Mi-Hua Tao, et al., 1993, J. Exp. Med). The antibody according to the present invention includes these antibodies having modified amino acid sequences in the constant regions. Modification of the amino acid sequence refers to deletion, substitution, or addition of one or several amino acids of the amino acid sequence. For example, the sequence CCC encoding proline (P) 331 in the constant region of the heavy chain according to the EU numbering system (see Sequences of proteins of immunological interest, NIH Publication No. 91-3242) is varied to TCC encoding serine (S) to substitute proline with serine, thereby changing the degree of binding to the complement. In the case of an anticancer agent, when the antibody itself does not have activity of inducing cell death, a preferable antibody has anti-tumor activity due to the antibody-dependent cellular cytotoxicity (ADCC) or complement-dependent cytotoxicity (CDC) through the Fc receptor. When the antibody itself has activity of inducing cell death, an antibody having a low degree of binding to the Fc receptor may be preferable. In the case of an immunosuppressive agent, an antibody having no ACDD or CDC activity is preferable when, for example, only the binding between the T-cell and the antigen-presenting cell is three-dimensionally suppressed. When ADCC or CDC activity could cause toxicity, an antibody having resolved the toxicity-causing activity by varying the Fc portion or changing the subclass may be preferable.

**[0041]** The antibody according to the present invention includes a modified antibody, the subclass of which has been rearranged to IgG1, IgG2, or IgG4, and a modified antibody, in which amino acid 331 in the constant region of the heavy chain of the modified antibody, the subclass of which has been rearranged to IgG1 or IgG2, according to the EU numbering system, has been substituted with Ser to constitute IgG1Ser or IgG2Ser. Specifically, the present invention includes an antibody in which only IgG1 and the IgG1Ser exhibit ADCC activity and only IgG1 and the IgG2 exhibit CDC activity as a result of the subclass modification into IgG1, IgG1Ser, IgG2, IgG2Ser, or IgG4.

**[0042]** Therapeutic effects on diseases such as cancer can be further enhanced by binding, for example, radionuclides such as iodine, yttrium, indium, and technetium (J. W. Goding, Monoclonal Antibodies: principles and practice, 1993, ACADEMIC PRESS), bacterial toxins such as *Pseudomonas exotoxin*, diphtheria toxin, and ricin, chemotherapeutics such as methotrexate, mitomycin, and calicheamicin (D. J. King, Applications and Engineering of Monoclonal Antibodies, 1998, T. J. International Ltd, M. L. Grossbard, Monoclonal Antibody-Based Therapy of Cancer, 1998, Marcel Dekker Inc), or a prodrug such as maytansinoid (Chari et al., Cancer Res., 1992, Vol. 52: 127, Liu et al., Proc Natl Acad Sci USA, 1996, Vol. 93: 8681) to the antibody according to the present invention.

**[0043]** In the present invention, the following steps are included in the production of a monoclonal antibody: (1) purification of a biopolymer that is used as an immunogen and/or production of a cell having excessively expressed antigen proteins on its surface; (2) production of an antibody-producing cell by immunizing an animal by injecting an antigen, sampling blood to assay the antibody titer, and determining the stage of excising the spleen or the like; (3) preparation of myeloma cells; (4) cell fusion between an antibody-producing cell and the myeloma cell; (5) selection of a group of hybridomas producing an antibody of interest; (6) division (cloning) into a single cell clone; (7) if necessary, culture of a hybridoma for mass-producing monoclonal antibodies or breeding animals to which the hybridoma has been transplanted; (8) examination of physiological activity or recognition specificity of the thus produced monoclonal antibody or examination of properties thereof as labeling reagents; and the like.

**[0044]** A process for producing the anti-HLA-DR monoclonal antibody is hereafter described in detail along with the above steps, although the process for producing the antibody is not limited thereto. For example, an antibody-producing cell other than a splenic or myeloma cell can also be used.

#### (1) Purification of antigen

**[0045]** A transformant is obtained by incorporating DNA encoding the HLA-DR  $\alpha$  chain and  $\beta$  chain into an expression vector for animal cells and introducing the expression vector into an animal cell. This can be used as an antigen in that state. Since the primary structures of the HLA-DR  $\alpha$  chain and  $\beta$  chain are known (Steven GE, et al., (2000), The HLA FactsBook, Academic Press), a peptide is chemically synthesized from the amino acid sequence of HLA-DR by a method known to a person skilled in the art, and the resultant can be used as an antigen.

**[0046]** A cell in which the full-length  $\alpha$  chain and  $\beta$  chain of the human HLA-DR are introduced into the L929 cell and HLA-DR heterodimers are excessively expressed on its surface is also effective as an immunogen. pEF-neo-HLA-DRA and pEF-neo-HLA-DR $\beta$  can be produced by separately incorporating DNA encoding the human HLA-DR  $\alpha$  chain protein and DNA encoding the human HLA-DR  $\beta$  chain protein into the expression vector for animal cells, pEF-neo. pEF-neo is a vector comprising a neomycin-resistant gene incorporated into modified pEF-BOS (see Mizushima S. & Nagata S., Nucleic Acids Res (1990), 18, 5332). It should be noted that DNA encoding HLA-DR, a vector, a host, or the like is not limited thereto.

[0047] Specifically, a transformant obtained by transforming the L929 cell in pEF-neo-HLA-DR $\alpha$  and pEF-neo-HLA-DR $\beta$  is cultured, and confirmation of HLA-DR expression using the trait of neomycin resistance acquired by the cell to which the pEF-neo vector has been inserted and the goat anti-HLA-DR polyclonal antibody (DAKO) are employed as indicators, thereby producing the L929 cell having excessively expressed human HLA-DR on its surface.

## (2) Step of preparing antibody-producing cell

[0048] The antigen obtained in (1) is mixed with an adjuvant such as Freund's complete or incomplete adjuvant or potash alum, and experimental animals are immunized with the resultant as the immunogen. A transgenic mouse capable of producing a human-derived antibody is most suitably used as an experimental animal, and such a mouse is described in literature by Tomizuka et al. (Tomizuka et al., Proc Natl Acad Sci USA, 2000, Vol. 97: 722).

[0049] An immunogen can be administered at the time of mouse immunization by any of hypodermic injection, intraperitoneal injection, intravenous injection, endodermic injection, intramuscular injection, or plantar injection. Intraperitoneal injection, plantar injection, or intravenous injection is preferable.

[0050] Immunization can be performed once or several times at suitable intervals (preferably at intervals of 2 to 4 weeks). Thereafter, the antibody titer against the antigen in the serum of the immunized animal is assayed, and the animal having a sufficiently high antibody titer is used as a source of antibody-producing cells. This can enhance the effects of the subsequent procedure. In general, the antibody-producing cell derived from an animal 3 to 5 days after the final immunization can be preferably used for the later cell fusion.

[0051] Examples of a method for measuring the antibody titer that is used herein include various conventional techniques such as radioimmunoassay (hereafter referred to as "RIA"), enzyme-linked immunosorbent assay (hereafter referred to as "ELISA"), fluorescent antibody technique, and passive haemagglutination. From the viewpoints of detection sensitivity, rapidity, accuracy, the possibility of automating operations, and the like, the use of RIA or ELISA is preferable.

[0052] In the present invention, the antibody titer can be assayed in the following manner in accordance with, for example, ELISA. At the outset, an antibody against a human antibody is allowed to adsorb on the surface of the solid phase such as a 96-well plate for ELISA. Subsequently, the surface of the solid phase having no antigen adsorbed thereon is covered with a protein that is unrelated to an antigen, for example, bovine serum albumin (BSA), the surface is washed, it is brought into contact with a gradually-diluted sample (for example, mouse serum) as a primary antibody, and the anti-HLA-DR antibody in the sample is then bound to the antigen. Further, an antibody against an enzyme-labeled human antibody is added as a secondary antibody and bound to a human antibody, followed by washing. Thereafter, the substrate of the enzyme is added, and changes of the absorbance, etc. are assayed based on the coloration due to substrate decomposition. Thus, the antibody titer is calculated.

## (3) Step of preparing myeloma cell

[0053] A cell incapable of producing autoantibodies derived from a mammalian such as a mouse, rat, guinea pig, hamster, rabbit, or human can be used as a myeloma cell. In general, established cell lines obtained from a mouse, for example, 8-azaguanine-resistant mouse (BALB/c-derived) myeloma cells P3X63Ag8U.1 (P3-U1) (Yelton, D.E. et al., Current Topics in Microbiology and Immunology, 81, 1-7 (1978)), P3/NSI/1-Ag4-1 (NS-1) (Kohler, G. et al. European J. Immunology, 6, 511-519 (1976)), Sp2/O-Ag14 (SP-2) (Shulman, M. et al. Nature, 276, 269-270 (1978)), P3X63Ag8.653 (653) (Kearney, J. F. et al. J. Immunology, 123, 1548-1550 (1979)), and P3X63Ag8 (X63) (Horibata, K. and Hams, A. W. Nature, 256, 495-497 (1975)) are preferably used. These cell lines are subjected to subculture in suitable medium, for example, 8-azaguanine medium (medium prepared by adding 8-azaguanine to RPMI-1640 medium comprising glutamine, 2-mercaptoethanol, gentamicin, and fetal calf serum (hereafter referred to as "FCS"), Iscove's Modified Dulbecco's Medium (hereafter referred to as "IMDM"), or Dulbecco's Modified Eagle Medium (hereafter referred to as "DMEM"). These cell lines are subjected to subculture in normal medium (for example, DMEM containing 10% FCS) 3 to 4 days before the cell fusion to reliably have  $2 \times 10^7$  or more cells on the day of the cell fusion.

## (4) Cell fusion

[0054] An antibody-producing cell is a plasma cell or a lymphocyte as its precursor cell. This may be obtained from any site of an individual and can be generally obtained from, for example, the spleen, lymph node, bone marrow, tonsilla, peripheral blood, or suitable combinations thereof, with the spleen cell being most commonly used.

[0055] After the final immunization, a site containing an antibody-producing cell therein, for example, the spleen, is excised from a mouse having a predetermined antibody titer to produce a spleen cell, which is an antibody-producing cell. The means for fusing this spleen cell with the myeloma cell obtained in step (3) that is most commonly performed at present is a method using polyethylene glycol having relatively low cytotoxicity and simple fusion operations. This



method comprises, for example, the following procedures.

**[0056]** Spleen cells and myeloma cells are thoroughly washed in serum-free medium (for example, DMEM) or phosphate-buffered saline (hereafter referred to as "PBS") and mixed with each other to bring the ratio of spleen cells to myeloma cells to approximately 5:1 to 10:1, followed by centrifugation. The supernatant is removed, the precipitated group of cells is thoroughly unraveled, and 1 mL of serum-free medium containing 50% (w/v) polyethylene glycol (molecular weight: 1,000 to 4,000) is then added thereto dropwise while stirring. Thereafter, 10 mL of serum-free medium is slowly added thereto, followed by centrifugation. The supernatant is discarded again, the precipitated cells are suspended in a suitable amount of normal medium containing a hypoxanthine-aminopterin-thymidine (hereafter referred to as "HAT") solution and human interleukin 6 (hereafter referred to as "IL-6") (this medium is hereafter referred to as "HAT medium"), the resultant is fractionated in each well of the culture plate (hereafter referred to as a "plate"), and cultured in the presence of 5% CO<sub>2</sub> at 37°C for approximately 2 weeks. During the culture, HAT medium is suitably supplemented.

#### (5) Selection of a group of hybridomas

**[0057]** When the aforementioned myeloma cells are 8-azaguanine-resistant, i.e., when they are hypoxanthine-guanine phosphoribosyltransferase (HGPRT) deficient, non-fused myeloma cells and cells fused between myeloma cells cannot survive in HAT-containing medium. While cells fused between antibody-producing cells or hybridomas of antibody-producing cells and myeloma cells can survive, the survival time of cells fused between antibody-producing cells is limited. Accordingly, continuation of culture in HAT-containing medium results in survival of only hybridomas of antibody-producing cells and myeloma cells. In consequence, this enables the selection of hybridomas.

**[0058]** HAT medium for the hybridomas grown as a colony is replaced with a medium from which aminopterin has been removed (hereafter referred to as "HT medium"). Thereafter, a part of the culture supernatant is collected to assay the anti-HLA-DR antibody titer by, for example, ELISA. When the aforementioned fusion protein is used as an antigen for ELISA, an operation of eliminating a clone is required so as not to select a clone that produces an antibody which specifically binds to the Fc region of the human IgG. The presence or absence of such a clone can be inspected by, for example, ELISA using the Fc region of the human IgG as an antigen.

**[0059]** A process using a 8-azaguanine-resistant cell line was exemplified above, although other cell lines can be also used depending on the process used for selecting a hybridoma. In such a case, the composition of the medium to be used is also changed.

#### (6) Step of cloning

**[0060]** The antibody titer is assayed in the same manner as described in (2), and the hybridomas, which were found to produce specific antibodies, are transferred to the other plate to perform cloning. Examples of cloning processes include: limiting dilution in which culture is conducted by diluting, so that one hybridoma is contained in a well of the plate; the soft agar method in which culture is conducted in a soft agar medium and colonies are recovered; a method in which culture is conducted by removing one cell using a micromanipulator; and a "sorter clone" process, in which one cell is separated using a cell sorter. Limiting dilution is simple and often employed.

**[0061]** The wells, the antibody titers of which have been recognized, are repeatedly subjected to cloning by, for example, limiting dilution 2 to 4 times, and those having stable antibody titers are selected as anti-HLA-DR monoclonal antibody-producing hybridomas.

**[0062]** The mouse-mouse hybridomas HD8, HD10, HD4, and HD6, the human anti-HLA-DR monoclonal antibody-producing cells according to the present invention, are deposited internationally at the International Patent Organism Depositary of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of October 11, 2001. The accession number of the hybridoma HD8 is FERM BP-7773, that of the hybridoma HD10 is FERM BP-7774, that of the hybridoma HD4 is FERM BP-7771, and that of the hybridoma HD6 is PERM BP-7772.

#### (7) Preparation of monoclonal antibody by culturing hybridomas

**[0063]** The hybridomas, which have completed the cloning process, are cultured in normal medium, which is a replacement of the HT medium. Large-scale culture is carried out by rotation culture in a large culture flask, spinner culture, or culture in a hollow-fiber system. The supernatant obtained through this large-scale culture is purified using a technique known to a person skilled in the art such as gel filtration. This enables the production of the anti-HLA-DR monoclonal antibody that comprises, as an active ingredient, the preventive or therapeutic agent of the present invention. Also, the hybridoma is multiplied in the abdominal cavity of a mouse of the same lineage (e.g., BALB/c), a nu/nu mouse, rat, guinea pig, hamster, or rabbit. This can provide ascites fluid containing a large amount of the anti-HLA-DR

monoclonal antibodies comprising, as an active ingredient, the preventive or therapeutic agent of the present invention. In order to simply carry out the purification, a commercially available monoclonal antibody purification kit (e.g., MabTrap GII Kit, Amersham Pharmacia Biotech) or the like can be used.

**[0064]** The thus obtained monoclonal antibody has high antigen specificity for the human HLA-DR.

#### (8) Examination of monoclonal antibody

**[0065]** The isotype and the subclass of the thus obtained monoclonal antibody can be identified in the following manner. Examples of a method for identification include the Ouchterlony method, ELISA, and RIA. Although the Ouchterlony method is simple, this method requires a concentrating operation when the concentration of the monoclonal antibody is low. In the case of ELISA or RIA, however, the isotype and the subclass of the monoclonal antibody can be identified by allowing the culture supernatant to react with the antigen-adsorbed solid phase, and using antibodies, as secondary antibodies, that correspond to various immunoglobulin isotypes and subclasses.

**[0066]** Proteins can be quantified by the Folin-Lowry method or by calculation based on the absorbance at 280 nm (1.4 (OD 280) = immunoglobulin 1 mg/ml).

**[0067]** The epitope recognized by the monoclonal antibody can be identified in the following manner. Various partial constructs of molecules recognized by the monoclonal antibody are first prepared. Partial constructs can be prepared by, for example, a method in which various partial peptides of the molecule are prepared using a conventional technique of synthesizing oligopeptides or a method in which a DNA sequence encoding a partial peptide of interest is incorporated in a suitable expression plasmid using a gene recombination technique to produce partial constructs inside or outside a host such as *E. coli*. These methods are generally used in combinations to attain the above object. For example, several polypeptide sequences, the lengths of which were successively and suitably shortened from the C-terminus or N-terminus of the antigen protein, are prepared by a gene recombination technique known to a person skilled in the art, and the reactivity of the monoclonal antibody therewith is then examined to roughly determine the recognition site.

**[0068]** Thereafter, various oligopeptides in the corresponding site, variants of the peptides, or the like are synthesized using a technique of synthesizing oligopeptides known to a person skilled in the art. The affinity of the monoclonal antibody, which is an active ingredient of the preventive or therapeutic agent of the present invention, with these peptides is then inspected or competition suppressing activity of peptides against the binding between the monoclonal antibody and the antigen is inspected, thereby more precisely limiting the epitope. In order to simply obtain various oligopeptides, a commercially available kit (for example, SPOTs Kit (Genosys Biotechnologies, Inc), a series of Multipin Peptide Synthesis Kits using the multipin synthesis technique (Chiron Corporation), or the like) can be used.

**[0069]** Alternatively, a gene encoding a human monoclonal antibody is cloned from an antibody-producing cell such as a hybridoma, the clone product is incorporated into a suitable vector, and the resultant is introduced into a host (e.g., a mammalian cell strain, *E. coli*, yeast cell, insect cell, or plant cell) to produce a recombinant antibody using a gene recombination technique (P. J. Delves, ANTIBODY PRODUCTION ESSENTIAL TECHNIQUES, 1997, WILEY, P. Shepherd and C. Dean, Monoclonal Antibodies, 2000 OXFORD UNIVERSITY PRESS, J. W. Goding, Monoclonal Antibodies: principles and practice, 1993, ACADEMIC PRESS).

**[0070]** The present invention includes a nucleic acid comprising a gene sequence of an antibody possessed by a hybridoma which produces the antibody of the present invention. More particularly, the present invention includes a nucleic acid that corresponds to the mature variable regions of the heavy chain and the light chain of the antibody produced from the hybridoma of the present invention as described below. The nucleic acid includes DNA and RNA. The present invention includes a nucleic acid, the sequence of which is modified by means of substitution, deletion, and/or addition of at least one nucleotide in the frame portion in the variable region of the heavy or light chain (see FR1, FR2, FR3, and FR4: Sequences of proteins of immunological interest, NIH Publication No. 91-3242), which hybridizes with a nucleic acid complementary to a nucleic acid before sequence modification under stringent conditions, binds to HLA-DR, has (a) life-extending effects in HLA-DR-expressed cancer cell-bearing non-human animals and (b) lower activity of suppressing immune responses compared with that of L243, and encodes an antibody exhibiting immunosuppressive activity equivalent to or higher than that of the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) in the nucleic acid comprising a gene sequence of the antibody possessed by a hybridoma which produces the antibody of the present invention. The antibody refers to an immunoglobulin in which a variable region and a constant region of the heavy chain and all regions including a variable region and a constant region of the light chain constituting the immunoglobulin are derived from a gene encoding the immunoglobulin. Stringent conditions involve the occurrence of hybridization only when the sequence is at least 90%, preferably at least 95%, and more preferably at least 97% homologous with the DNA sequence encoding the antibody of the present invention. In general, these conditions involve the occurrence of hybridization at a temperature about 5°C to about 30°C, and preferably about 10°C to about 25°C lower than the melting temperature of the perfect hybrid. Stringent conditions are described in J. Sambrook et al., Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press (1989), and the conditions described therein can be used.

[0071] In order to prepare a gene encoding the monoclonal antibody from a hybridoma, DNAs encoding the V region of the L chain, the C region of the L chain, the V region of the H chain, and the C region of the H chain of the monoclonal antibody are prepared by PCR and the like. Oligo DNA constructed from the anti-HLA-DR antibody gene or the amino acid sequence can be used as a primer, and DNA prepared from a hybridoma can be used as a template. These DNAs are incorporated into a suitable vector, and the resultant is introduced into a host to be expressed. Alternatively, these DNAs are separately incorporated into each suitable vector for co-expression.

[0072] A phage or plasmid that can autonomously multiply in a host microorganism is used as a vector. Examples of plasmid DNA include a plasmid derived from *E. coli*, *Bacillus subtilis*, or yeast, and an example of phage DNA is  $\lambda$  phage.

[0073] A host that is used in transformation is not particularly limited as long as the gene of interest can be expressed therein. Examples thereof include bacteria (e.g., *E. coli* or *Bacillus subtilis*), yeast, animal cells (e.g., COS cells or CHO cells), and insect cells.

[0074] A method for introducing a gene into a host is known, and examples thereof include any methods such as a method using calcium ions, electroporation, spheroplast, the lithium acetate method, the calcium phosphate method, and lipofection. Examples of a method for introducing a gene into animals as described below include microinjection, a method for introducing a gene into the ES cell by electroporation or lipofection, and nucleus transplantation.

[0075] In the present invention, the anti-HLA-DR antibody can be obtained by collecting it from a culture product obtained by culturing a transformant. The "culture product" refers to any of: (a) a culture supernatant; (b) a cultured cell, cultured bacterial cell, or fragmented product thereof; or (c) a secretion product of the transformant. When culturing a transformant, a medium that is suitable for the host of interest is used, and a culture method such as stationary culture or roller bottle culture is employed.

[0076] After the culture, when the protein of interest is produced in a bacterial or other cell, the bacterial or other cell is destroyed to collect the antibody. When the antibody of interest is produced outside the bacterial or other cell, the culture solution remaining unchanged is used, or the bacterial or other cell is removed by centrifugation, etc. Thereafter, general biochemical techniques using various types of chromatography for isolation and purification of proteins are performed singly or in suitable combinations. Thus, the antibody of interest can be isolated and purified from the culture product.

[0077] With the use of a technique for preparing transgenic animals, animal hosts comprising the genes of the antibody of interest are incorporated in endogenous genes. For example, transgenic cattle, goats, sheep, or pigs are prepared. From the milk secreted from these transgenic animals, a large amount of monoclonal antibodies derived from the antibody genes thereof can be obtained (Wright, G. et al., (1991), *Bio/Technology* 9, 830-834). When culturing a hybridoma *in vitro*, a hybridoma is multiplied, maintained, and stored in accordance with various conditions such as properties of the cells to be cultured, purposes of tests and research, or culture methods. A known nutrient medium that is used for producing a monoclonal antibody in a culture supernatant or various nutrient media derived and prepared from a known basal medium can be used to perform the culture.

#### (9) Properties of the antibody

[0078] The antibody according to the present invention has the following functional properties a) and b), and these properties can be confirmed by, for example, methods described for each item:

a) HLA-DR-expressing human cancer cells are transplanted in immunodeficient mice such as SCID mice, the survival ratio of mice when inoculated with the antibody of the present invention is inspected, and as a result, the number of days for which the mice survive is prolonged; and

b) activity of suppressing immune responses by allogeneic mixed lymphocyte reaction is lower than that of L243. More specifically, the properties a) and b) are as follows:

(a) when a 6-week-old SCID mouse is inoculated intravenously with 10  $\mu$ L of the anti-asialo GM1 antiserum, on the next day, inoculated intravenously with 5 x 10<sup>6</sup> of Burkitt's lymphoma cells Raji (ATCC CCL-86), and 5 days thereafter, inoculated once intravenously with 5 to 50  $\mu$ g/kg, and preferably 5  $\mu$ g/kg, based on body weight, of the antibody of the present invention, the survival ratio of the mouse 90 days after the inoculation is higher than that 90 days after the inoculation with the same amount of the human anti-HSA antibody; and  
(b) the immunosuppressive activity is lower than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) at the same concentration, wherein the immunosuppressive activity is assayed as follows: 50  $\mu$ L of antibody adjusted at 8 to 200 mg/mL, and preferably 8 mg/mL using 10% FCS-containing RPMI 1640 medium is mixed with 50  $\mu$ L of mature dendritic cell suspension derived from a first human donor adjusted at 2 x 10<sup>5</sup> cells/mL using 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate, the mixture is allowed to stand at 4°C for 30 minutes, the resultant is mixed with 100  $\mu$ L of T-cell suspension

(purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using 10% FCS-containing RPMI 1640 medium derived from a second human donor having a histocompatible antigen different from that of the first human donor, the mixture is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 5 days, <sup>3</sup>H thymidine is added theret at 1.0 μCi/well, the resultant is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 16 to 20 hours, the <sup>3</sup>H thymidine incorporated in the cell is recovered and then measured using a scintillator, and the incorporation of the <sup>3</sup>H thymidine into the cell is used as an indicator to assay the immunosuppressive activity.

**[0079]** Examples of such an antibody include an antibody produced from the hybridoma HD (accession number: FERM BP-7771), an antibody produced from the hybridoma HD8 (accession number: FERM BP-7773), and an antibody produced from the hybridoma HD10 (accession number FERM BP-7774).

**[0080]** The aforementioned property a) indicates that the antibody has potent anti-tumor activity.

**[0081]** The antibody according to the present invention has a functional property, that is, activity of suppressing immune responses by allogeneic mixed lymphocyte reaction is equivalent to or higher than that of L243. More specifically, the immunosuppressive activity is equivalent to or higher than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55 at the same concentration, wherein the immunosuppressive activity is assayed as follows. First, 50 μl of antibody adjusted at 8-200 mg/mL, preferably 8 mg/mL using 10% FCS-containing RPM 1640 medium is mixed with 50 μL of mature dendritic cell suspension derived from a first human donor adjusted at  $2 \times 10^5$  cells/mL using 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate. The mixture is then allowed to stand at 4°C for 30 minutes, and the resultant is mixed with 100 μL of T-cell suspension (purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using 10% FCS-containing RPMI 1640 medium derived from a second human donor having ; histocompatible antigen different from that of the first human donor. The resultant is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 5 days, <sup>3</sup>H thymidine is added thereto at 1.0 μCi/well, the resultant is further cultured at 37°C in the presence of 5% CO<sub>2</sub> for 16 to 20 hours, the <sup>3</sup>H thymidine incorporated in the cell is recovered and then measured using a scintillator, and incorporation of the <sup>3</sup>H thymidine into the cell is used as an indicator to assay the immunosuppressive activity. An example of such an antibody is one produced from the hybridoma HD6 (accession number: FERM BP-7772).

**[0082]** The antibody having the aforementioned activity according to the present invention is useful as an ingredient for a preventive or therapeutic agent for malignant tumors or an ingredient for an immunosuppressive agent.

**[0083]** Even more surprisingly, the antibody according to the present invention significantly suppress a lowering in the survival ratio of tumor-bearing mouse models caused by the tumor cell growth at a low dose of 0.1 μg per mouse (5 μg per kg of body weight), and exhibits life extending-effects in mouse models. If the survival ratio of the mice to which the antibody of the present invention has been administered is significantly enhanced compared with that of the control mice when the human anti-human serum albumin (HSA) antibody is administered as a control simultaneously with the antibody according to the present invention, the antibody of the present invention can be determined to exhibit the life extending-effects. For example, the antibody according to the present invention and the anti-HSA antibody are administered to 5 each of tumor-bearing mouse models to which lymphoma cells have been transplanted. If at least one mouse to which the antibody of the present invention has been administered is alive when all of the mice in the group of mice to which the anti-HSA antibody has been administered have died, it can be said that the life extending-effects can be exhibited in tumor-bearing mouse models.

**[0084]** The immunosuppressive effects can be evaluated based on activity of suppressing the immune response by allogeneic mixed lymphocyte reaction (MLR) as described above, and MLR can be carried out in a conventional manner.

**[0085]** Also, the epitope of the HLA-DR recognized by the antibody of the present invention can be identified by a conventional method. For example, regarding 199 amino acids in the extracellular region of the HLA-DR β chain (DRB1\*15011) (199 amino acids from amino acids 29 to 227 in the amino acid sequence as shown in SEQ ID NO: 147, and the nucleotide sequence encoding the amino acid sequence as shown in SEQ ID NO: 147 is shown in SEQ ID NO: 146), peptides are prepared by shifting an amino acid in the 13-mer peptides (for example, peptides having amino acid sequences as shown in SEQ ID NOs: 52 to 145), and the reactivity is inspected. In such a case, the antibody according to the present invention maximally binds to the peptide having the amino acid sequence as shown in SEQ ID NO: 82, or potentially binds to at least one of the three peptides as shown in SEQ ID NOs: 82, 83, and 84. The term "potentially binds" refers to the binding which exhibits fluorescence intensity at least 10 times as great as that of the background in the method as described in Example 17 (2). The antibody of the present invention has reactivity with peptides 61 to 71 of the HLA-DR β chain. Further, HLA-DR is known to have approximately 350 types of polymorphisms (see the IMGT/HLA database of EMBL-EBI, etc.). The antibody according to the present invention is capable of recognizing substantially almost all of these polymorphisms. Whether or not this is true can be determined by preparing a group of peptides including substantially almost all of the approximately 350 types of polymorphisms and assaying the reactivity with substantially almost all of these peptides. For example, if an antibody reacts with 12 or more types of peptides among 16 types of peptides as shown in SEQ ID NOs: 24 to 39, this antibody is capable of recognizing substantially almost all of the polymorphisms of the HLA-DR. The antibody according to the present invention signifi-

cantly binds to all the peptides as shown in SEQ ID NOs: 24 to 39 and all the peptides as shown in SEQ ID NOs: 40 to 43. The term "significantly binds" refers to the binding which exhibits fluorescence intensity at least 10% as great as the background in the method as described in Example 17 (3).

## 2. Pharmaceutical composition

**[0086]** A preparation that comprises a purified preparation of the human anti-HLA-DR antibody of the present invention is also within the scope of the present invention. Such a preparation preferably comprises a physiologically acceptable diluent or carrier in addition to the antibody, and it may be a mixture of the aforementioned antibody with another antibody or another agent such as an antibiotic. Examples of a suitable carrier include, but are not limited to, physiological saline, phosphate-buffered saline, phosphate-buffered saline glucose solution, and a buffered saline solution. Alternatively, the antibody may be lyophilized and used by being recomposed with the addition of the above buffered aqueous solution, when it is needed. The preventive or therapeutic agent can be administered in various dosage forms, and examples of dosage forms include oral administration in the form of, for example, a tablet, capsule, granule, powder, or syrup and parenteral administration in the form of, for example, an injection, drop, or suppository.

**[0087]** The dose may vary depending on symptom, age, body weight, etc. In the case of oral administration, the dose is generally about 0.01 mg to 1,000 mg per day per adult, and this amount of preparation can be administered in one dose or several separate doses. In the case of parenteral administration, about 0.01 mg to 1,000 mg can be administered per dose through hypodermic injection, intramuscular injection, or intravenous injection.

**[0088]** The antibody or a pharmaceutical composition according to the present invention can be applied to the treatment or prevention of various diseases or symptoms that could be caused by an HLA-DR-expressing cell. Examples of such diseases or symptoms include various malignant tumors, and examples of the application thereof are use as an immunosuppressive agent at the time of organ transplantation (a preventive or therapeutic agent for immunological rejection at the time of pancreatic islet or kidney transplantation, or GVHD), a therapeutic agent for autoimmune diseases (for example, rheumatism, arteriosclerosis, multiple sclerosis, systemic erythematodes, idiopathic thrombocytopenia, or Crohn's disease), and a therapeutic agent for allergic diseases such as asthma.

**[0089]** For example, the antibody according to the present invention and a pharmaceutical composition comprising this antibody having the following functional properties a) and b) can be used for preventing or treating various malignant tumors.

a) HLA-DR-expressing human cancer cells are transplanted in immunodeficient mice such as SCID mice, the survival ratio of mice when inoculated with the antibody of the present invention is inspected, and as a result, the number of days for which the mice survive is prolonged; and

b) activity of suppressing immune responses by allogeneic mixed lymphocyte reaction is lower than that of L243.

**[0090]** The antibody according to the present invention and a pharmaceutical composition comprising this antibody having activity of suppressing immune responses by allogeneic mixed lymphocyte reaction equivalent to or higher than that of L243 can be used for preventing or treating rheumatism or graft-versus-host disease (GvHD).

**[0091]** The present invention includes a process for preventing or treating the aforementioned diseases using the antibody or pharmaceutical composition according to the present invention. The present invention also includes the use of the antibody according to the present invention in the production of a preventive or therapeutic agent for the aforementioned diseases.

**[0092]** Examples of tumors that can be prevented or treated are leukemia (including chronic lymphatic leukemia and acute lymphatic leukemia), lymphoma (including non-Hodgkin's lymphoma, Hodgkin's lymphoma, T-cell lymphoma, B-cell lymphoma, Burkitt's lymphoma, malignant lymphoma, diffuse lymphoma, and follicular lymphoma), myeloma (including multiple myeloma), breast cancer, colon cancer, kidney cancer, gastric cancer, ovarian cancer, pancreatic cancer, cervical cancer, endometrial cancer, esophageal cancer, liver cancer, head and neck squamous cancer, skin cancer, urinary tract cancer, prostate cancer, choriocarcinoma, pharyngeal cancer, laryngeal cancer, pleural tumor, arrhenoblastoma, endometrial hyperplasia, endometriosis, embryoma, fibrosarcoma, Kaposi's sarcoma, angioma, cavernous angioma, hemangioblastoma, retinoblastoma, spongiocytoma, neurofibroma, oligodendroglioma, medulloblastoma, neuroblastoma, neuroglioma, rhabdomyoblastoma, glioblastoma, osteogenic sarcoma, leiomyosarcoma, thyroid sarcoma, and Wilms tumor. When the antibody of the present invention is applied, the kind of a tumor is not limited to one, and combinations of several kinds of tumors may be involved.

## 3. Preparation example

**[0093]** The molecule according to the present invention is used as an ampule for an aseptic solution or suspension dissolved in water or another pharmaceutically acceptable solution. Also, the ampule is filled with an aseptic powder

preparation (preferably lyophilized molecules of the present invention), and it may be diluted with a pharmaceutically acceptable solution at the time of use thereof.

#### Brief Description of the Drawings

[0094]

Fig. 1 is a diagram showing the immunosuppressive activity of the purified human anti-HLA-DR monoclonal antibody on MLR.

Fig. 2 is a diagram showing the life-extending effects of the purified human anti-HLA-DR monoclonal antibody on the tumor-bearing mouse model. Fig. 2A shows a case where the dose is 1 µg/head, and Fig. 2B shows the effects attained when the dose is 0.1 µg/head.

Fig. 3 is a diagram showing the antibody-dependent cellular cytotoxicity (ADCC) and the complement-dependent cytotoxicity (CDC) of the purified human anti-HLA-DR monoclonal antibody. Fig. 3A shows the ADCC activities of HD8G1Ser and HD8G1, Fig. 3B shows the CDC activities of HD8G2CHO, HD8G1, HD8G1Ser, and HD8G4, Fig. 3C shows the ADCC activities of HD8G1, HD8G2, HD8G2Ser, HD8G4, HD4G1, HD4G2Ser, and HD4G4, and Fig. 3D shows the CDC activities of HD8G1, HD8G2, HD8G2Ser, HD8G4, HD4G1, HD4G2Ser, and HD4G4.

Fig. 4 is a diagram showing the life-extending effects of the antibodies according to the present invention, HD8G1Ser, HD8G2Ser, and HD8G1, on the tumor-bearing mouse model. Fig. 4A shows the results attained when the dose is 0.1 µg per mouse, and Fig. 4B shows the results attained when the dose is 1.0 µg per mouse.

Fig. 5 is a diagram showing the analysis of the purified human anti-HLA-DR monoclonal antibody epitope using the synthetic peptide in the "LA-DR β chain sequence. Figs. 5A, 5B, and 5C show analyses on HD4, HD6, and HD8, respectively.

Fig. 6 is a diagram showing the analysis of the purified human anti-HLA-DR monoclonal antibody epitope using the synthetic peptide in the polymorphic sequence in the HLA-DR β chain epitope sequence.

#### Free text of Sequence Listing

[0095]

SEQ ID Nos: 1 to 15: description of artificial sequence: primer

SEQ ID Nos: 24 to 145: description of artificial sequence: peptide

#### Best Modes for Carrying out the Invention

[0096] The present invention is hereafter described in more detail with reference to the following examples, although the technical scope of the present invention is not limited to the embodiments described in these examples.

#### Example 1: Preparation of antigen

[0097] In order to obtain a cell having excessively expressed human HLA-DR on its cell membrane, plasmid vectors for the expression of full-length amino acids in the human HLA-DR α chain and β chain were prepared. DNA encoding the HLA-DR α chain and β chain was prepared by PCR.

#### a) Preparation of expression vectors for full-length human HLA-DR α chain and β chain

[0098] In template PCR, plasmid vectors pEF-neo-HLA-DRα and pEF-neo-HLA-DRβ holding cDNA encoding the human HLA-DR α chain and β chain were used as the templates. pEF-neo-HLA-DRα and pEF-neo-HLA-DRβ were prepared in the manner as described below. DNA of the full-length human HLA-DR α chain and that of the HLA-DR β chain were modified by the polymerase chain reaction (PCR) for adding the EcoRI sequence at the 5'-terminus and the NotI sequence and a stop codon at the 3'-terminus. PCR was carried out for 30 PCR cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 60 seconds using, as a template, the cDNA derived from human peripheral blood mononuclear cells, as primers, synthesizing 5'-CCGGAATTCACCATGGCCATAAGTGGAGTCCCTGTG -3' (SEQ ID NO: 1) with 5'-AAAGCGGCCGCTCATTACAGAGGCCCTGCGTTCTGC -3' (SEQ ID NO: 2) for the HLA-DRα and synthesizing 5'-CCGGAATTCCTGGTCTGCTGTTCTCCAGCA-3' (SEQ ID NO: 3) with 5'-AAAGCGGCCGCTCATCAGCTCAGGAATCCTGTTGGCTG -3' (SEQ ID NO: 4) for the HLA-DRβ, and the LA-Taq DNA polymerase (Gibco BRL). The synthesized sequence was isolated as the EcoRI-NotI fragment and linked to the pEF-neo vector cleaved with the same enzyme (a vector comprising the neomycin-resistant gene incorporated in modified

pEF-BOS (see Mizushima S. & Nagata S., Nucleic Acids Res (1990), 18, 5332)). The resulting plasmids were designated as pEF-neo-HLA-DR $\alpha$  and pEF-neo-HLA-DR $\beta$ . 765 bp cDNA was encoded in the HLA-DR $\alpha$  incorporated in pEF-neo-HLA-DR $\alpha$ , and 801 bp cDNA was encoded in the HLA-DR $\beta$  incorporated in pEF-neo-HLA-DR $\beta$ . In all of PCRs in the following examples, the reaction temperature was regulated using the Gene Amp PCR System 9700 (Perkin Elmer Japan).

#### c) Preparation of a human HLA-DR-expressing cell

**[0099]** pEF-neo-HLA-DR $\alpha$  and pEF-neo-HLA-DR $\beta$  prepared in b) were introduced in the L929 cell (American Type Culture Collection (ATCC) No. CCL-1) using LipofectAMINE Plus (Gibco BRL). Gene introduction was carried out in accordance with the method described in the instructions therefor. The gene-introduced cells were cultured in a cell culture flask (culture area: 75 cm<sup>2</sup>) at 37°C in the presence of 5.0% CO<sub>2</sub> for 24 hours, and G418 (Gibco BRL) was added thereto at 1 mg/mL, followed by culture for a week. Subsequently, flow cytometry (FCM, Becton Dickinson) was carried out using the R-phycoerythrin-labeled mouse anti-human-HLA-DR antibody (BD Pharmingen), and cells having HLA-DR expressed on the surfaces of their cell membranes were selectively sorted from those that had attained G418-resistance among the gene-introduced cells.

**[0100]** All oligonucleotides such as PCR primers were synthesized using a DNA automatic synthesizer (Model: 3948, Perkin-Elmer's Applied Biosystems Division) in accordance with the attached instructions (see Matteucci, M.D. and Caruthers, M.H., (1981), J. Am. Chem. Soc. 103, 3185-3191). After the completion of the synthesis, each of the oligonucleotides was cleaved from the support and then deprotected. The resulting solution was exsiccated, dissolved in distilled water, and then cryopreserved at -20°C before use.

#### Example 2: Preparation of human antibody-producing mouse

**[0101]** The mouse used for immunization is genetically homozygous for both endogenous Ig heavy chain breakdown and  $\kappa$  light chain breakdown, and simultaneously retains a chromosome 14 fragment (SC 20) containing a human Ig heavy chain locus and a human Ig $\kappa$  chain transgene (KCoS). This mouse was prepared by mating a mouse having a human Ig heavy chain locus (lineage A) with a mouse having a human Ig $\kappa$  chain transgene (lineage B). Lineage A is homozygous for both endogenous Ig heavy chain breakdown and  $\kappa$  light chain breakdown, and retains a chromosome 14 fragment (SC20) that is transmittable. For example, it is described in the report by Tomizuka et al. (Tomizuka, et al., Proc. Natl. Acad. Sci. USA, 2000, Vol. 97: 722). Lineage B is homozygous for both endogenous Ig heavy chain breakdown and  $\kappa$  light chain breakdown, and retains a human Ig $\kappa$  chain transgene (KCo5, a transgenic mouse). It is described for example, in the report by Fishwild et al. (Nat. Biotechnol., 1996, Vol. 14: 845).

**[0102]** An individual obtained by mating a male mouse of lineage A with a female mouse of lineage B or a female mouse of lineage A with a male mouse of lineage B in which the human Ig heavy chain and the  $\kappa$  light chain can be simultaneously detected in the serum (Ishida & Lonberg, IBC's 11th Antibody Engineering, Abstract 2000) was used in the following immunity experiment. The aforementioned human antibody-producing mouse (referred to as a "KM mouse") can be obtained from Kirin Brewery Co., Ltd. by making a contract.

#### Example 3: Preparation of human monoclonal antibody against human HLA-DR

**[0103]** In this example, a monoclonal antibody was prepared in accordance with a common technique as described in, for example, "*Tan-kuron koutai jikken sousa nyuumon* (A guide to monoclonal antibody experiments)" (Tamie ANDO et al., Kodansha Ltd. Publishers, 1991). The HLA-DR-expressing L929 cell prepared in Example 1 was used as the immunogen human HLA-DR. A human immunoglobulin-producing human antibody-producing mouse prepared in Example 2 was used as an animal to be immunized.

**[0104]** In order to prepare a human monoclonal antibody against human HLA-DR, human antibody-producing mice were subjected to initial immunization intraperitoneally with the HLA-DR-expressing L929 cells (5 x 10<sup>6</sup> cells per mouse) prepared in Example 1. After the initial immunization, the same cells were administered for immunization 2, 4, and 8 weeks later. Three days before obtaining the spleen and the lymph node as described below, cells (1 x 10<sup>6</sup> cells per mouse) were administered to the tail veins, and recombinant human IL-6 (hereafter referred to as "IL-6," 5 ng per mouse, prepared at the Pharmaceutical Research Laboratory, Kirin Brewery Co., Ltd.) was administered subcutaneously.

**[0105]** The spleen and/or lymph node were surgically obtained from the immunized mouse, the recovered organs were placed in 10 mL of serum-free DMEM medium containing 350 mg/mL sodium bicarbonate, 50 units/mL of penicillin, and 50  $\mu$ g/mL of streptomycin (Gibco BRL, hereafter referred to as "serum-free DMEM medium"), and mashed on a mesh (cell strainer, Falcon) using a spatula. The cell suspension that had passed through the mesh was centrifuged to precipitate the cells. Thereafter, these cells were washed twice in serum-free DMEM medium and then suspended

in serum-free DMEM medium to count the number of cells. In contrast, myeloma cells SP2/0 (ATCC No. CRL-1581), which were cultured in DMEM medium (Gibco BRL) containing 10% FCS (SIGMA) (this medium is hereafter referred to as "serum-containing DMEM medium") at 37°C in the presence of 5% CO<sub>2</sub> in such a manner that the cell concentration did not exceed 1 x 10<sup>6</sup> cells/mL, were similarly washed in serum-free DMEM medium and then suspended in serum-free DMEM medium to count the number of cells. The recovered cell suspension and the mouse myeloma cell suspension were mixed with each other at a ratio of 5:1, the mixture was centrifuged, and the supernatant was completely removed. 1 mL of 50% (w/v) polyethylene glycol 1500 (Boehringer Mannheim) was slowly added as a fusing agent to the pellet while stirring the pellet with the tip of a pipette, 1 mL of serum-free DMEM medium previously heated at 37°C was slowly added two separate times, and 7 mL of serum-free DMEM medium was further added. After the centrifugation, the supernatant was removed, and the resulting fusion cell was subjected to screening by limiting dilution as described below. A hybridoma was selected by culturing it in DMEM medium containing 10% FCS, IL-6 (10 ng/mL), and hypoxanthine (H), aminopterin (A), and thymidine (T) (hereafter referred to as "HAT," SIGMA). Further, a single clone was obtained by limiting dilution using HT (SIGMA), 10% FCS, and IL-6-containing DMEM medium. Culture was conducted in a 96-well microtiter plate (Becton Dickinson). The selection (screening) of a hybridoma clone that produces the anti-human HLA-DR human monoclonal antibody and the characterization of the human monoclonal antibody that is produced by each hybridoma were carried out with enzyme-linked immunosorbent assay (ELISA) and FCM as described later.

**[0106]** In the screening of human monoclonal antibody-producing hybridomas, many human monoclonal antibody-producing hybridomas were obtained, which have human immunoglobulin  $\gamma$  chain (hig $\gamma$ ) and human immunoglobulin light chain  $\kappa$ , and which have specific reactivity with human HLA-DR through Cell ELISA as described in Examples 4 and 5.

Example 4: Selection of human anti-HLA-DR monoclonal antibody-producing clones having human immunoglobulin light chain  $\kappa$  (Ig $\kappa$ )

**[0107]** Burkitt's lymphoma cells, Daudi (ATCC No. CCL-213), were added to each well in quantities of 1 x 10<sup>5</sup> cells, the hybridoma supernatant was added thereto, and the mixture was incubated at 4°C for 20 minutes. Subsequently, the incubation product was washed twice with 1 % FCS-containing PBS, the horseradish peroxidase-labeled goat anti-human immunoglobulin light chain  $\kappa$  (Ig $\kappa$ ) antibody (50  $\mu$ g/well, DAKO) was added thereto, and the resultant was incubated at 4°C for 20 minutes. The incubation product was washed twice with 1% FCS-containing PBS, and 100  $\mu$ L each of a TMB chromogenic substrate solution (DAKO) was added to each well, followed by incubation at room temperature for 20 minutes. 0.5M sulfuric acid (100  $\mu$ L/well) was added to each well to terminate the reaction. Absorbance at a 450 nm wavelength (reference wavelength: 570 nm) was assayed using a microplate reader (1420 ARVO multilabel counter, Wallac) to select positive antibody-producing clones.

Example 5: Identification of subclass of each monoclonal antibody

**[0108]** After the addition of 1 x 10<sup>5</sup> Daudi cells to each well, the hybridoma supernatant was added thereto, and the resultant was incubated at 4°C for 20 minutes. Subsequently, the incubation product was washed twice with 1 % FCS-containing PBS, and horseradish peroxidase-labeled sheep anti-human IgG1 antibody, sheep anti-human IgG2 antibody, sheep anti-human IgG3 antibody, or sheep anti-human IgG4 antibody (2000-fold diluted, 50  $\mu$ L/well, The Binding Site) was added to each well, followed by incubation at room temperature for 1 hour. After being washed three times with 1% FCS-containing PBS, a substrate buffer (TMB, 100  $\mu$ L/well, DAKO) was added to each well, and incubation was carried out at room temperature for 20 minutes. Subsequently, 0.5M sulfuric acid (100  $\mu$ L/well) was added to terminate the reaction. Absorbance at a 450 nm wavelength (reference wavelength: 570 nm) was assayed using a microplate reader (1420 ARVO multilabel counter, Wallac) to identify the subclass for each clone. Clones which were not positive for any subclass were excluded from the selection since they were not IgG

**[0109]** The results only on the finally selected clones are shown in Table 1.

Table 1

Properties of selected anti-HLA-DR monoclonal antibodies				
	Subclass	Activity of suppressing immune responses by allogeneic mixed lymphocyte reaction	Reactivity with L929	Reactivity with L929/HLA-DR
Human IgG	Poly	-	-	-
HD3	IgG1	++	-	+



Table 1 (continued)

Properties of selected anti-HLA-DR monoclonal antibodies				
	Subclass	Activity of suppressing immune responses by allogeneic mixed lymphocyte reaction	Reactivity with L929	Reactivity with L929/HLA-DR
HD4	IgG1	++	-	+
HD6	IgG1	+++	-	+
HD7	IgG3	+	-	+
HD8	IgG2	+	-	+
HD10	IgG2	+	-	+

Example 6: Process for obtaining normal human mononuclear cell and normal human dendritic cell

**[0110]** At the outset, a normal human peripheral blood-derived mononuclear cell was prepared in accordance with a conventional method using Ficoll (Ficoll-PaquePLUS, Amersham Pharmacia Biotech). The normal human blood contained in a blood-sampling bag (Terumo) containing a sodium citrate solution as an anticoagulant was centrifuged (600 G, room temperature, 5 minutes) to separate a cell fraction from blood plasma. The cell fraction was diluted with PBS, superposed on Ficoll, and mononuclear cells were separated by specific gravity-based centrifugation (400 G, room temperature, 30 minutes). The intermediate layer was extracted as a mononuclear cell and washed twice with PBS. The resultant was further diluted with PBS, and centrifuged at 100 G for 10 minutes to remove blood platelets remaining in the supernatant. Thus, normal human peripheral blood-derived mononuclear cells (PBMC) were obtained.

**[0111]** Subsequently, the obtained PBMC was allowed to react with CD14 antibody attached to magnetic beads (Miltenyi Biotec (MB)) at 4°C for 30 minutes, and positive selection for CD14-positive cells was carried out through a MACS separating column (MB). The MACS separating column was used in accordance with the attached instruction. The CD14-positive cells were cultured in 10% FCS-containing RPMI medium comprising GM-CSF (final concentration: 50 ng/mL, prepared at the Pharmaceutical Research Laboratory, Kirin Brewery Co., Ltd.) and interleukin 4 (final concentration: 200 ng/mL, Genzyme Corporation) for 5 to 8 days. Thereafter, lipopolysaccharide (LPS, final concentration: 40 ng/mL, Difco) was added thereto, and the resultant was cultured overnight. The culture product was then used as a mature dendritic cell (mature DC).

Example 7: Activity of suppressing immune responses by allogeneic mixed lymphocyte reaction

**[0112]** In an allogeneic transplantation having different major histocompatibility antigens (MHC), the T-cell is activated by recognizing the nonself (histoincompatible) MHC molecular complex (alloantigen), thereby generating immunological rejection. The human MHC is referred to as human leukocyte antigen (HLA), and there are class I antigens to which HLA-A, B, and C belong, and class II antigens to which HLA-DP, DQ, and DR belong. Further, since each molecule has a polymorphic property, several thousand combinations of human HLA are possible. Thus, histoincompatibility is very highly likely to occur with another person. The allogeneic mixed lymphocyte culture is a test to inspect *in vitro* the growth of T-cells which react with alloantigens by subjecting lymphocytes having different histocompatible antigens (hereafter referred to as donor A and donor B for convenience) to mixed culture.

**[0113]** Activity of suppressing the immune response by allogeneic mixed lymphocytes was assayed using the culture supernatant of hybridomas selected in Examples 4 and 5. A normal human peripheral blood mononuclear cell was obtained in the manner as described in Example 6. The mononuclear cells of donor A were allowed to react in RPMI 1640 medium containing mitomycin C (25 µg/mL, 37°C, 30 minutes) to suppress the multiplication of cells. After the reaction, they were washed at least three times in RPMI 1640 medium and suspended in RPMI 1640 medium at  $1 \times 10^6$ /mL. The mononuclear cells of donor B were fractionated in a 96-well plate at  $1 \times 10^5$  cell/well, the medium was removed by centrifugation, the culture supernatant was added at 100 µL/well, and the resultant was allowed to stand at 4°C for 30 minutes.

**[0114]** Subsequently, the mononuclear cells of donor A were fractionated at 100 µL/well in a 96-well plate containing the mononuclear cells of donor B and the culture supernatant, followed by culturing at 37°C for 4 days in the presence of 5% CO<sub>2</sub>. Thereafter, <sup>3</sup>H thymidine (Amersham Pharmacia Biotech) was added thereto at 1.0 µCi/well, and culture was further conducted at 37°C for 16 to 20 hours in the presence of 5% CO<sub>2</sub>. The <sup>3</sup>H thymidine incorporated in cells were collected on a glass filter mat (Printed Filtermat, Wallac) using the Micro96 Harvester (SKATRON), dehydrated, well immersed in scintillator (Betap, Scint, Wallac), and packaged. Thereafter, activity of the β dose was assayed using a liquid scintillation counter (1205 BETAPLATE, Wallac).

**[0115]** Properties of the anti-human HLA-DR antibodies selected as a result of the assay are shown in Table 1.

Among those, HD8, HD10, HD4, and HD6 are deposited internationally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6,1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of October 11, 2001. The accession number of the hybridoma HD8 is FERM BP-7773, that of the hybridoma HD10 is FERM BP-7774, that of the hybridoma HD4 is FERM BP-7771, and that of the hybridoma HD6 is FERM BP-7772.

#### Example 8: Preparation of each antibody

**[0116]** Thu human anti-HLA-DR monoclonal antibody was purified from the culture supernatant of hybridomas obtained in Examples 4 and 5 in the following manner. The culture supernatant containing the human anti-HLA-DR monoclonal antibody was subjected to affinity purification using rmp Protein A (Amersham Pharmacia Biotech) and a 0.8 x 4.0 cm column (Biorad), PBS as an adsorption buffer, and a 0.02M glycine buffer (pH 3) as an elution buffer. The elution fraction was adjusted at around pH 7.2 with the addition of 1M Tris (pH 9.0). The prepared antibody solution was converted into PBS using a dialysis membrane (molecular weight cutoff: 10,000, Spectrum Laboratories), filter-sterilized with a membrane filter MILLEX-GV (pore diameter: 0.22  $\mu$ m, MILLIPORE), and the purified human anti-HLA-DR monoclonal antibody was obtained. The concentration of the purified antibody was calculated by assaying the absorbance at 280 nm and on the basis of 1 mg/mL = 1.4 OD.

**[0117]** The culture supernatant containing the human anti-HLA-DR monoclonal antibody was prepared in the following manner. At the outset, the human anti-HLA-DR monoclonal antibody-producing hybridoma was adapted to eRDF medium (Kyokuto Seiyaku) containing 10 ng/ml of IL-6 and 10% fetal calf serum (FCS, SIGMA). Subsequently, a part thereof was adapted to eRDF medium (Kyokuto Seiyaku) containing bovine insulin (5  $\mu$ g/ml, Gibco BRL), human transferin (5  $\mu$ g/ml, Gibco BRL), ethanolamine (0.01 mM, SIGMA), sodium selenite ( $2.5 \times 10^{-5}$  mM, SIGMA), and 1% low IgG FCS (HyClone) to purify the antibody. These adapted hybridomas were cryopreserved. Culture was conducted in a flask, and the culture supernatant was collected when the rate of surviving hybridomas reached 90%. The collected supernatant was applied to 10  $\mu$ m- and 0.2  $\mu$ m-filters (Gelman Science) to eliminate waste such as hybridomas.

#### Example 9: Activity of suppressing immune responses by allogeneic mixed lymphocytes by purified human anti-HLA-DR monoclonal antibody

**[0118]** The allogeneic mixed lymphocyte culture is a test to inspect *in vitro* the growth of T-cells which react with alloantigens by subjecting lymphocytes having different histocompatible antigens (hereafter referred to as donor A and donor B for convenience) to mixed culture. The mature dendritic cell (DC) derived from a monocyte *in vitro* induced from the peripheral blood of donor A and the T-cells separated from the peripheral blood of donor B are exclusively subjected to mixed culture. This enables the inspection of similar reactivity of T-cells with an alloantigen when the antibody-dependent cellular cytotoxicity (ADCC) of the macrophage, etc. has been removed. Mixed culture using allogeneic DC and T-cells was carried out in the presence of the anti-HLA-DR monoclonal antibody to inspect the function of the anti-HLA-DR monoclonal antibody on the T-cell alloantigen reactivity. Specifically, the mature DC of donor A was suspended in 10% FCS-containing RPMI 1640 medium (RPMI-10% FCS) at  $2 \times 10^5$  cells/mL on a round-bottom, 96-well plate (Falcon). The anti-HLA-DR monoclonal antibody, the human polyclonal IgG (hIgG) as a negative control, and the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) as a positive control were diluted with RPMI-10% FCS to 200, 40, and 8 mg/mL, respectively. The T-cells separated from the peripheral blood of donor B (purity: 99% or higher) were suspended in RPMI-10% FCS at  $1 \times 10^6$  cells/mL. At the outset, 50  $\mu$ L of the mature DC derived from donor A was mixed with the same amount of the anti-HLA-DR monoclonal antibody on a 96-well plate, and the resultant was allowed to stand at 4°C for 30 minutes. Subsequently, 100  $\mu$ L of T-cells derived from donor B was mixed therewith, cultured at 37°C for 5 days in the presence of 5% CO<sub>2</sub>, <sup>3</sup>H thymidine (Amersham Pharmacia Biotech) was added at 1.0  $\mu$ Ci/well, and culture was further conducted at 37°C for 16 to 20 hours in the presence of 5% CO<sub>2</sub>. The <sup>3</sup>H thymidine incorporated in the cells was collected on a glass filter mat (Printed Filtermat, Wallac) using the Micro96 Harvester (SKATRON), dehydrated, well immersed in a scintillator (Betap, Scint, Wallac), and packaged. Thereafter, activity of the  $\beta$ -ray dose was assayed using a liquid scintillation counter (1205 BETAPLATE, Wallac).

**[0119]** The results are shown in Fig. 1. The immunosuppressive activities of HD4, HD6, HD8, and HD10 were more dose-dependent compared with those of the hIgG group, those of HD8, HD4, and HD10 were lower than those of L243, and those of HD6 were equivalent to or higher than those of the positive control antibody L243.

**[0120]** This suggests that HD8, HD4, and HD10 can be used as antibodies having low immunosuppressive activities, and HD6 can be used as an immunosuppressive agent.

#### Example 10: Examination of reactivity of each monoclonal antibody with HLA-DR-expressing cells

**[0121]** Reactivity of each purified monoclonal antibody obtained in Example 3 with the HLA-DR-expressing L929

cells prepared in Example 1 was analyzed by FCM. The L929 cells and the HLA-DR-expressing L929 cells were suspended in PBS containing 0.1% sodium azide and 1% fetal calf serum (Staining Medium, hereinafter abbreviated to "SM") at  $2 \times 10^7$ /mL, and the suspension was fractionated in a 96-well, round-bottom plate at 100  $\mu$ L/well. After the centrifugation (600 G, 4°C, 2 minutes), the supernatant was removed, the culture supernatant (50  $\mu$ L) of hybridomas cultured in Example 3 was added, and the mixture was stirred and allowed to stand under ice cooling for 30 minutes. Centrifugation (600 G, 4°C, 2 minutes) was then carried out to remove the supernatant. The pellet was washed twice with 100  $\mu$ L/well SM, 30  $\mu$ L of 0.0125 mg/mL RPE fluorescence-labeled rabbit anti-human Ig $\kappa$  F(ab')<sub>2</sub> antibody (DAKO) was added thereto, and the resultant was incubated under ice cooling for 30 minutes. After being washed twice with SM, the incubation product was suspended in SM, and the fluorescence intensity of each cell was assayed by FCM.

[0122] The results are shown in Table 1 above. All the antibodies exhibited potent binding activities to HLA-DR-expressing L929 cells alone, but none exhibited the binding activity to the L929 cell. This indicates that these antibodies specifically bind to HLA-DR.

#### Example 11: Effect of purified human anti-HLA-DR monoclonal antibody on tumor-bearing mouse models

[0123] The effects of the purified human anti-HLA-DR monoclonal antibody obtained in Example 8 were examined using tumor-bearing mouse models in accordance with a method described below.

[0124] First, 5-week-old C.B-17/ICR SCID mice (CLEA Japan, Inc.) were purchased, anti-asialo GM1 antiserum (Wako Chemicals) was diluted, and 10  $\mu$ L each thereof was intravenously administered to each of the mice when they were 6 weeks old. On the next day, Burkitt's lymphoma cells Raji (ATCC CCL-86) were intravenously administered in amounts of  $5 \times 10^6$  cells per mouse. Five days after the Raji transplantation, the purified human anti-HLA-DR monoclonal antibody was administered once in tail veins of mice in amounts of 0.1  $\mu$ g or 1  $\mu$ g per mouse. As an antibody negative control, the same amount of human anti-HSA antibody was used. The survival ratio after the transplantation was observed for about 3 months or longer.

[0125] The results of the experiment are shown in Fig. 2. In the group of mice to which the purified human anti-HLA-DR monoclonal antibody was administered in amounts of 0.1  $\mu$ g per mouse, the number of surviving mice 90 days later was 3 for HD8, and 1 for HD10 and HD4 out of a group of 5 mice (Fig. 2B). In contrast, in the group of mice to which the antibody was administered in amounts of 1.0  $\mu$ g per mouse, the number of surviving mice was 3 for HD8, 5 for HD10, and 2 for HD 4 out of a group of 5 mice (Fig. 2A). In the negative control group to which the anti-HSA antibody was administered, all mice died within 60 days after the Raji transplantation.

[0126] At the time of the antibody administration, the body weight of each mouse was about 20 g. Thus, 0.1  $\mu$ g and 1  $\mu$ g per mouse means 5  $\mu$ g/kg and 50  $\mu$ g/kg based on body weight, respectively. This revealed that HD8 exhibits an anti-tumor effect at very low dosages. Based on this and the results attained in Example 9 together, HD8 can be said to be an antibody having low immunosuppressive activity and potent anti-tumor activity. This suggests that HD8 can be used as an anti-cancer agent with few side effects.

#### Example 12: Preparation of a gene encoding a monoclonal antibody and construction of a recombinant antibody-expressing vector

##### (1) cDNA cloning of HD4 and HD8 antibody genes and preparation of expression vector

[0127] Hybridomas HD4 and HD8 were cultured in eRDF medium (Kyokuto Seiyaku) containing 10 ng/ml of IL-6 (R & D Systems) and 10% fetal bovine serum (SIGMA) and centrifuged to collect cells. Thereafter, TRIZOL (Gibco BRL) was added thereto, and total RNA was extracted in accordance with the instructions therefor. Cloning of the variable region in the antibody cDNA was carried out using the SMART RACE cDNA amplification Kit (Clontech) in accordance with the attached instruction.

[0128] First-strand cDNA was prepared using 5  $\mu$ g of the total RNA as a template.

##### 1) Synthesis of first-strand cDNA

[0129] Total RNA 5  $\mu$ g/3  $\mu$ L

5'CDS 1  $\mu$ L

SMART oligo 1  $\mu$ L

[0130] The reaction solution having the above composition was incubated at 70°C for 2 minutes,

5 $\times$  Buffer 2 $\mu$ L

DTT 1  $\mu$ L

DNTP mix 1  $\mu$ L

Superscript II 1  $\mu$ L

were then added, and the mixture was incubated at 42°C for 1.5 hours.

**[0131]** Further, 100 µL of Tricine Buffer was added, followed by incubation at 72°C for 7 minutes. Thus, first-strand cDNA was obtained.

- 5 2) Amplification of a heavy chain gene and a light chain gene by PCR and construction of a recombinant antibody-expressing vector

**[0132]** cDNA was amplified using Z-Taq (Takara).

cDNA 2 µL

10 10xZ-Taq Buffer 5 µL

dNTP mix 4 µL

Z-Taq 1 µL

Primer 1

Primer 2

15 **[0133]** The final volume of the reaction solution having the above composition was brought to 50 µL with the aid of double distilled water and then subjected to PCR.

**[0134]** The heavy chain was amplified by 30 PCR cycles of 98°C for 1 second and 68°C for 30 seconds using UMP (SMART RACE cDNA amplification Kit, Clontech) and the hh-6 primer (5'-GGT CCG GGA GAT CAT GAG GGT GTC CTT-3') (SEQ ID NO: 5). Further, 1 µL of this reaction solution was used as a template, and 20 PCR cycles of 98°C for 1 second and 68°C for 30 seconds were repeated using NUMP (SMART RACE cDNA amplification Kit, Clontech) and the hh-3 primer (5'-GTG CAC GCC GCT GGT CAG GGC GCC TG-3') (SEQ ID NO: 6). Thereafter, the amplified PCR product was purified using a PCR purification kit (QIAGEN), and nucleotide sequencing was carried out using hh-4 (5'-GGT GCC AGG GGG AAG ACC GAT GG-3') (SEQ ID NO: 7) as a primer. Based on the sequence information, the HD4 heavy chain specific primer tnHD4Sal (5'-ata tgt cga cCC AGC CCT GGG ATT TTC AGG TGT TTT C-3') (SEQ ID NO: 8) and the HD8 heavy chain specific primer tnHD8Sal (5'-ata tgt cga cTGG CTG ACC AGG GCA GTC ACC AGA G-3') (SEQ ID NO: 9) were synthesized, and the resultant primer was used to determine the sequence also from the opposite direction. PCR was carried out using a specific primer and tnCHNhe (5'-gat ggg ccc ttg gtg cta get gag gag acg g-3') (SEQ ID NO: 10) (98°C for 1 second, 60°C for 30 seconds, and 72°C for 30 seconds), the amplified cDNA fragment of the heavy chain was digested with Sail and NheI, and introduced into the NSKG1-Val Lark vector (a modified vector of IDEC Pharmaceuticals, N5KG1 (US patent 6,001,358)) that was cleaved with the same enzyme. The inserted sequence was confirmed to be identical to the one identified by direct sequencing by identifying the sequence using the vector as a template.

**[0135]** The light chain was amplified by repeating 30 PCR cycles of 98°C for 1 second and 68°C for 30 seconds using UMP (SMART RACE cDNA amplification Kit, Clontech) and the hk-2 primer (5'-GTT GAA GCT CTT TGT GAC GGG CGA GC -3') (SEQ ID NO: 11). Further, 1 µL of this reaction solution was used as a template, and 20 PCR cycles of 98°C for 1 second and 68°C for 30 seconds were repeated using NUMP (SMART RACE cDNA amplification Kit, Clontech) and hk-6 (5'-TGCG GGG AAG ATG AAG ACA GAT GGT G-3') (SEQ ID NO: 12). Thereafter, the amplified PCR product was purified using a PCR purification kit (QIAGEN), and nucleotide sequencing was carried out using the hk-6 primer (SEQ ID NO: 12). Based on the sequence information, the HD4 light chain specific primer tnHD4Bgl (5'-ata tag atc tGC TGC TCA GTT AGG ACC CAG AGG GAA CC-3') (SEQ ID NO: 13) and the HD8 light chain specific primer tnHD8Bgl (5'-ata tag atc tGG GAG TCA GAC CCA CTC AGG ACA CAG C-3') (SEQ ID NO: 14) were synthesized, and the resultant primer was used to determine the sequence also from the opposite direction. PCR was carried out using a specific primer and tnCkBsi (5'-aag aca gat ggt gca gcc acc gta cgt ttg at-3') (SEQ ID NO: 15) (98°C for 1 second, 60°C for 30 seconds, and 72°C for 30 seconds), the amplified cDNA fragment of the light chain was digested with BglII and BsiWI and introduced into the N5KG1-Val Lark vector that was cleaved with the same enzyme. The inserted sequence was confirmed to be identical to the one identified by direct sequencing by identifying the sequence using the vector as a template.

**[0136]** DNAs encoding the variable region of the heavy chain, that of the light chain of HD4, amino acid sequences of the variable region of the heavy chain, and that of the light chain of HD4 are shown below.

50

55

<The variable region of the heavy chain of HD4> (SEQ ID NO: 16)

[0137]

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55  
 60  
 65  
 70  
 75  
 80  
 85  
 90  
 95  
 100  
 105  
 110  
 115  
 120  
 125  
 130  
 135  
 140  
 145  
 150  
 155  
 160  
 165  
 170  
 175  
 180  
 185  
 190  
 195  
 200  
 205  
 210  
 215  
 220  
 225  
 230  
 235  
 240  
 245  
 250  
 255  
 260  
 265  
 270  
 275  
 280  
 285  
 290  
 295  
 300  
 305  
 310  
 315  
 320  
 325  
 330  
 335  
 340  
 345  
 350  
 355  
 360  
 365  
 370  
 375  
 380  
 385  
 390  
 395  
 400  
 405  
 410  
 415  
 420  
 425  
 430  
 435  
 440  
 445  
 450  
 455  
 460  
 465  
 470  
 475  
 480  
 485  
 490  
 495  
 500  
 505  
 510  
 515  
 520  
 525  
 530  
 535  
 540  
 545  
 550  
 555  
 560  
 565  
 570  
 575  
 580  
 585  
 590  
 595  
 600  
 605  
 610  
 615  
 620  
 625  
 630  
 635  
 640  
 645  
 650  
 655  
 660  
 665  
 670  
 675  
 680  
 685  
 690  
 695  
 700  
 705  
 710  
 715  
 720  
 725  
 730  
 735  
 740  
 745  
 750  
 755  
 760  
 765  
 770  
 775  
 780  
 785  
 790  
 795  
 800  
 805  
 810  
 815  
 820  
 825  
 830  
 835  
 840  
 845  
 850  
 855  
 860  
 865  
 870  
 875  
 880  
 885  
 890  
 895  
 900  
 905  
 910  
 915  
 920  
 925  
 930  
 935  
 940  
 945  
 950  
 955  
 960  
 965  
 970  
 975  
 980  
 985  
 990  
 995  
 1000  
 1005  
 1010  
 1015  
 1020  
 1025  
 1030  
 1035  
 1040  
 1045  
 1050  
 1055  
 1060  
 1065  
 1070  
 1075  
 1080  
 1085  
 1090  
 1095  
 1100  
 1105  
 1110  
 1115  
 1120  
 1125  
 1130  
 1135  
 1140  
 1145  
 1150  
 1155  
 1160  
 1165  
 1170  
 1175  
 1180  
 1185  
 1190  
 1195  
 1200  
 1205  
 1210  
 1215  
 1220  
 1225  
 1230  
 1235  
 1240  
 1245  
 1250  
 1255  
 1260  
 1265  
 1270  
 1275  
 1280  
 1285  
 1290  
 1295  
 1300  
 1305  
 1310  
 1315  
 1320  
 1325  
 1330  
 1335  
 1340  
 1345  
 1350  
 1355  
 1360  
 1365  
 1370  
 1375  
 1380  
 1385  
 1390  
 1395  
 1400  
 1405  
 1410  
 1415  
 1420  
 1425  
 1430  
 1435  
 1440  
 1445  
 1450  
 1455  
 1460  
 1465  
 1470  
 1475  
 1480  
 1485  
 1490  
 1495  
 1500  
 1505  
 1510  
 1515  
 1520  
 1525  
 1530  
 1535  
 1540  
 1545  
 1550  
 1555  
 1560  
 1565  
 1570  
 1575  
 1580  
 1585  
 1590  
 1595  
 1600  
 1605  
 1610  
 1615  
 1620  
 1625  
 1630  
 1635  
 1640  
 1645  
 1650  
 1655  
 1660  
 1665  
 1670  
 1675  
 1680  
 1685  
 1690  
 1695  
 1700  
 1705  
 1710  
 1715  
 1720  
 1725  
 1730  
 1735  
 1740  
 1745  
 1750  
 1755  
 1760  
 1765  
 1770  
 1775  
 1780  
 1785  
 1790  
 1795  
 1800  
 1805  
 1810  
 1815  
 1820  
 1825  
 1830  
 1835  
 1840  
 1845  
 1850  
 1855  
 1860  
 1865  
 1870  
 1875  
 1880  
 1885  
 1890  
 1895  
 1900  
 1905  
 1910  
 1915  
 1920  
 1925  
 1930  
 1935  
 1940  
 1945  
 1950  
 1955  
 1960  
 1965  
 1970  
 1975  
 1980  
 1985  
 1990  
 1995  
 2000  
 2005  
 2010  
 2015  
 2020  
 2025  
 2030  
 2035  
 2040  
 2045  
 2050  
 2055  
 2060  
 2065  
 2070  
 2075  
 2080  
 2085  
 2090  
 2095  
 2100  
 2105  
 2110  
 2115  
 2120  
 2125  
 2130  
 2135  
 2140  
 2145  
 2150  
 2155  
 2160  
 2165  
 2170  
 2175  
 2180  
 2185  
 2190  
 2195  
 2200  
 2205  
 2210  
 2215  
 2220  
 2225  
 2230  
 2235  
 2240  
 2245  
 2250  
 2255  
 2260  
 2265  
 2270  
 2275  
 2280  
 2285  
 2290  
 2295  
 2300  
 2305  
 2310  
 2315  
 2320  
 2325  
 2330  
 2335  
 2340  
 2345  
 2350  
 2355  
 2360  
 2365  
 2370  
 2375  
 2380  
 2385  
 2390  
 2395  
 2400  
 2405  
 2410  
 2415  
 2420  
 2425  
 2430  
 2435  
 2440  
 2445  
 2450  
 2455  
 2460  
 2465  
 2470  
 2475  
 2480  
 2485  
 2490  
 2495  
 2500  
 2505  
 2510  
 2515  
 2520  
 2525  
 2530  
 2535  
 2540  
 2545  
 2550  
 2555  
 2560  
 2565  
 2570  
 2575  
 2580  
 2585  
 2590  
 2595  
 2600  
 2605  
 2610  
 2615  
 2620  
 2625  
 2630  
 2635  
 2640  
 2645  
 2650  
 2655  
 2660  
 2665  
 2670  
 2675  
 2680  
 2685  
 2690  
 2695  
 2700  
 2705  
 2710  
 2715  
 2720  
 2725  
 2730  
 2735  
 2740  
 2745  
 2750  
 2755  
 2760  
 2765  
 2770  
 2775  
 2780  
 2785  
 2790  
 2795  
 2800  
 2805  
 2810  
 2815  
 2820  
 2825  
 2830  
 2835  
 2840  
 2845  
 2850  
 2855  
 2860  
 2865  
 2870  
 2875  
 2880  
 2885  
 2890  
 2895  
 2900  
 2905  
 2910  
 2915  
 2920  
 2925  
 2930  
 2935  
 2940  
 2945  
 2950  
 2955  
 2960  
 2965  
 2970  
 2975  
 2980  
 2985  
 2990  
 2995  
 3000  
 3005  
 3010  
 3015  
 3020  
 3025  
 3030  
 3035  
 3040  
 3045  
 3050  
 3055  
 3060  
 3065  
 3070  
 3075  
 3080  
 3085  
 3090  
 3095  
 3100  
 3105  
 3110  
 3115  
 3120  
 3125  
 3130  
 3135  
 3140  
 3145  
 3150  
 3155  
 3160  
 3165  
 3170  
 3175  
 3180  
 3185  
 3190  
 3195  
 3200  
 3205  
 3210  
 3215  
 3220  
 3225  
 3230  
 3235  
 3240  
 3245  
 3250  
 3255  
 3260  
 3265  
 3270  
 3275  
 3280  
 3285  
 3290  
 3295  
 3300  
 3305  
 3310  
 3315  
 3320  
 3325  
 3330  
 3335  
 3340  
 3345  
 3350  
 3355  
 3360  
 3365  
 3370  
 3375  
 3380  
 3385  
 3390  
 3395  
 3400  
 3405  
 3410  
 3415  
 3420  
 3425  
 3430  
 3435  
 3440  
 3445  
 3450  
 3455  
 3460  
 3465  
 3470  
 3475  
 3480  
 3485  
 3490  
 3495  
 3500  
 3505  
 3510  
 3515  
 3520  
 3525  
 3530  
 3535  
 3540  
 3545  
 3550  
 3555  
 3560  
 3565  
 3570  
 3575  
 3580  
 3585  
 3590  
 3595  
 3600  
 3605  
 3610  
 3615  
 3620  
 3625  
 3630  
 3635  
 3640  
 3645  
 3650  
 3655  
 3660  
 3665  
 3670  
 3675  
 3680  
 3685  
 3690  
 3695  
 3700  
 3705  
 3710  
 3715  
 3720  
 3725  
 3730  
 3735  
 3740  
 3745  
 3750  
 3755  
 3760  
 3765  
 3770  
 3775  
 3780  
 3785  
 3790  
 3795  
 3800  
 3805  
 3810  
 3815  
 3820  
 3825  
 3830  
 3835  
 3840  
 3845  
 3850  
 3855  
 3860  
 3865  
 3870  
 3875  
 3880  
 3885  
 3890  
 3895  
 3900  
 3905  
 3910  
 3915  
 3920  
 3925  
 3930  
 3935  
 3940  
 3945  
 3950  
 3955  
 3960  
 3965  
 3970  
 3975  
 3980  
 3985  
 3990  
 3995  
 4000  
 4005  
 4010  
 4015  
 4020  
 4025  
 4030  
 4035  
 4040  
 4045  
 4050  
 4055  
 4060  
 4065  
 4070  
 4075  
 4080  
 4085  
 4090  
 4095  
 4100  
 4105  
 4110  
 4115  
 4120  
 4125  
 4130  
 4135  
 4140  
 4145  
 4150  
 4155  
 4160  
 4165  
 4170  
 4175  
 4180  
 4185  
 4190  
 4195  
 4200  
 4205  
 4210  
 4215  
 4220  
 4225  
 4230  
 4235  
 4240  
 4245  
 4250  
 4255  
 4260  
 4265  
 4270  
 4275  
 4280  
 4285  
 4290  
 4295  
 4300  
 4305  
 4310  
 4315  
 4320  
 4325  
 4330  
 4335  
 4340  
 4345  
 4350  
 4355  
 4360  
 4365  
 4370  
 4375  
 4380  
 4385  
 4390  
 4395  
 4400  
 4405  
 4410  
 4415  
 4420  
 4425  
 4430  
 4435  
 4440  
 4445  
 4450  
 4455  
 4460  
 4465  
 4470  
 4475  
 4480  
 4485  
 4490  
 4495  
 4500  
 4505  
 4510  
 4515  
 4520  
 4525  
 4530  
 4535  
 4540  
 4545  
 4550  
 4555  
 4560  
 4565  
 4570  
 4575  
 4580  
 4585  
 4590  
 4595  
 4600  
 4605  
 4610  
 4615  
 4620  
 4625  
 4630  
 4635  
 4640  
 4645  
 4650  
 4655  
 4660  
 4665  
 4670  
 4675  
 4680  
 4685  
 4690  
 4695  
 4700  
 4705  
 4710  
 4715  
 4720  
 4725  
 4730  
 4735  
 4740  
 4745  
 4750  
 4755  
 4760  
 4765  
 4770  
 4775  
 4780  
 4785  
 4790  
 4795  
 4800  
 4805  
 4810  
 4815  
 4820  
 4825  
 4830  
 4835  
 4840  
 4845  
 4850  
 4855  
 4860  
 4865  
 4870  
 4875  
 4880  
 4885  
 4890  
 4895  
 4900  
 4905  
 4910  
 4915  
 4920  
 4925  
 4930  
 4935  
 4940  
 4945  
 4950  
 4955  
 4960  
 4965  
 4970  
 4975  
 4980  
 4985  
 4990  
 4995  
 5000  
 5005  
 5010  
 5015  
 5020  
 5025  
 5030  
 5035  
 5040  
 5045  
 5050  
 5055  
 5060  
 5065  
 5070  
 5075  
 5080  
 5085  
 5090  
 5095  
 5100  
 5105  
 5110  
 5115  
 5120  
 5125  
 5130  
 5135  
 5140  
 5145  
 5150  
 5155  
 5160  
 5165  
 5170  
 5175  
 5180  
 5185  
 5190  
 5195  
 5200  
 5205  
 5210  
 5215  
 5220  
 5225  
 5230  
 5235  
 5240  
 5245  
 5250  
 5255  
 5260  
 5265  
 5270  
 5275  
 5280  
 5285  
 5290  
 5295  
 5300  
 5305  
 5310  
 5315  
 5320  
 5325  
 5330  
 5335  
 5340  
 5345  
 5350  
 5355  
 5360  
 5365  
 5370  
 5375  
 5380  
 5385  
 5390  
 5395  
 5400  
 5405  
 5410  
 5415  
 5420  
 5425  
 5430  
 5435  
 5440  
 5445  
 5450  
 5455  
 5460  
 5465  
 5470  
 5475  
 5480  
 5485  
 5490  
 5495  
 5500  
 5505  
 5510  
 5515  
 5520  
 5525  
 5530  
 5535  
 5540  
 5545  
 5550  
 5555  
 5560  
 5565  
 5570  
 5575  
 5580  
 5585  
 5590  
 5595  
 5600  
 5605  
 5610  
 5615  
 5620  
 5625  
 5630  
 5635  
 5640  
 5645  
 5650  
 5655  
 5660  
 5665  
 5670  
 5675  
 5680  
 5685  
 5690  
 5695  
 5700  
 5705  
 5710  
 5715  
 5720  
 5725  
 5730  
 5735  
 5740  
 5745  
 5750  
 5755  
 5760  
 5765  
 5770  
 5775  
 5780  
 5785  
 5790  
 5795  
 5800  
 5805  
 5810  
 5815  
 5820  
 5825  
 5830  
 5835  
 5840  
 5845  
 5850  
 5855  
 5860  
 5865  
 5870  
 5875  
 5880  
 5885  
 5890  
 5895  
 5900  
 5905  
 5910  
 5915  
 5920  
 5925  
 5930  
 5935  
 5940  
 5945  
 5950  
 5955  
 5960  
 5965  
 5970  
 5975  
 5980  
 5985  
 5990  
 5995  
 6000  
 6005  
 6010  
 6015  
 6020  
 6025  
 6030  
 6035  
 6040  
 6045  
 6050  
 6055  
 6060  
 6065  
 6070  
 6075  
 6080  
 6085  
 6090  
 6095  
 6100  
 6105  
 6110  
 6115  
 6120  
 6125  
 6130  
 6135  
 6140  
 6145  
 6150  
 6155  
 6160  
 6165  
 6170  
 6175  
 6180  
 6185  
 6190  
 6195  
 6200  
 6205  
 6210  
 6215  
 6220  
 6225  
 6230  
 6235  
 6240  
 6245  
 6250  
 6255  
 6260  
 6265  
 6270  
 6275  
 6280  
 6285  
 6290  
 6295  
 6300  
 6305  
 6310  
 6315  
 6320  
 6325  
 6330  
 6335  
 6340  
 6345  
 6350  
 6355  
 6360  
 6365  
 6370  
 6375  
 6380  
 6385  
 6390  
 6395  
 6400  
 6405  
 6410  
 6415  
 6420  
 6425  
 6430  
 6435  
 6440  
 6445  
 6450  
 6455  
 6460  
 6465  
 6470  
 6475  
 6480  
 6485  
 6490  
 6495  
 6500  
 6505  
 6510  
 6515  
 6520  
 6525  
 6530  
 6535  
 6540  
 6545  
 6550  
 6555  
 6560  
 6565  
 6570  
 6575  
 6580  
 6585  
 6590  
 6595  
 6600  
 6605  
 6610  
 6615  
 6620  
 6625  
 6630  
 6635  
 6640  
 6645  
 6650  
 6655  
 6660  
 6665  
 6670  
 6675  
 6680  
 6685  
 6690  
 6695  
 6700  
 6705  
 6710  
 6715  
 6720  
 6725  
 6730  
 6735  
 6740  
 6745  
 6750  
 6755  
 6760  
 6765  
 6770  
 6775  
 6780  
 6785  
 6790  
 6795  
 6800  
 6805  
 6810  
 6815  
 6820  
 6825  
 6830  
 6835  
 6840  
 6845  
 6850  
 6855  
 6860  
 6865  
 6870  
 6875  
 6880  
 6885  
 6890  
 6895  
 6900  
 6905  
 6910

<The variable region of the light chain of HD4> (SEQ ID NO: 19)

[0140]

METPAQLLFLLLLWLPDTTGELVLTSQSPGTLSPGERATLSCRASQSVSSRYLAWYQQK  
PGQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPLTFGGG  
TKVEIKRT

[0141] The translation initiation site in the heavy chain DNA is the ATG codon starting from the 79th adenine (A) from the 5'-terminus of the sequence as shown in SEQ ID NO: 16, and the boundary between the variable region and the constant region of the antibody is located between the 504th adenine (A) and the 505th guanine (G) from the 5'-terminus. In the amino acid sequence, the variable region of the heavy chain is a portion between the N-terminus and the 142nd serine (S) residue in the sequence as shown in SEQ ID NO: 17, and a portion comprising the 143rd alanine (A) and thereafter is the constant region. The N-terminus of the purified heavy chain protein was analyzed. This demonstrated that the signal sequence of the heavy chain was a portion between the N-terminus and the 19th cysteine (C) of the sequence as shown in SEQ ID NO: 17, and the N-terminus of the matured body was the 20th glutamic acid (E) of the sequence as shown in SEQ ID NO: 17. Accordingly, the matured portion in the amino acid sequence as shown in SEQ ID NO: 17 is a portion between the 20th glutamic acid and the 142nd serine.

[0142] The translation initiation site in the light chain DNA is the ATG codon starting from the 35th A from the 5'-terminus of the sequence as shown in SEQ ID NO: 18, and the variable region is the portion between the 5'-terminus and the 418th adenine (A). In the amino acid sequence, the variable region is the portion between the N-terminus and the 128th lysine (K) of the sequence as shown in SEQ ID NO: 19. The N-terminus of the purified light chain protein was analyzed. This demonstrated that the signal sequence of the light chain was the portion between the N-terminus and the 20th glycine (G) of the sequence as shown in SEQ ID NO: 19, and the N-terminus of the matured body was the 21st glutamic acid (E) of the sequence as shown in SEQ ID NO: 19. Accordingly, the matured portion in the amino acid sequence as shown in SEQ ID NO: 20 is the portion between the 21st glutamic acid and the 128th lysine.

[0143] DNAs encoding the variable region of the heavy chain, that of the light chain of HD8, amino acid sequences of the variable region of the heavy chain, and that of the light chain of HD8 are shown below.

<The variable region of the heavy chain of HD8> (SEQ ID NO: 20)

[0144]

GTCGACTGGCTGACCAGGGCAGTCACCAGAGCTCCAGACAATGTCTGTCTCCTTCCTC  
ATCTTCCTGCCCCGTGCTGGGCCTCCCATGGGGTGTCTGTACAGGTTTCAGCTGCAGC  
ACTCAGGTCCAGGACTGGTGAAGCCCTCGCAGACCCTCTCACTCACCTGTGCCATCTC  
CGGGGACAGTGTCTCTAGCAACAGTGCTTCTTGGAACTGGATCAGGCAGTCCCCATCG  
AGGGGCCTTGAGTGGCTGGGAAGGACATACTACAGGTCCAAGTGGTATAATGATTATG  
CAGTATCTGTGAAAAGTCGAATAGTCATCAACCCAGACACATCCAAGAACCAGTTCTC  
CCTGCAGCTGAACTCTGTGACTCCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAA  
AATTTCTATGGTTCCGGAGACTTGTGATAAGAAGTATTACTGCTACGGTATGGACGTCTG  
GGGCCAAGGGACCACGGTCACCGTCTCCTCAGCTAGC

<The variable region of the heavy chain of HD8> (SEQ ID NO: 21)

[0145]

MSVSFLIFLPVLGLPWGVLSQVQLQHSGPGLVKPSQTL~~SLT~~CAISGDSVSSNSASWNWIRQ  
SPSRGLEWLGRYYRSKWYNDYAVSVKSRVINPDTSKNQFSLQLNSVTPEDTAVYYCAR  
ENFYGSETCHKKYYCYGMDVWVGQGTTVTVSSAS

<The variable region of the light chain of HD8> (SEQ ID NO: 22)

[0146]

AGATCTGGGAGTCAGACCCACTCAGGACACAGCATGGACATGAGGGTCCCCGCTCAG  
CTCCTGGGGGCTTCTGCTGCTCTGGCTCCCAGGTGCCAGATGTGCCATCCAGTTGACCC  
AGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGC  
AAGTCAGGGCATTAGCAGTGCTTTAGCCTGGTATCAGCAGAAACCAGGGAAAGCTCCT  
AAACTCCTGATCTATGATGCCTCCAGTTTGGAAGTGGGGTCCCATCAAGGTTACGCG  
GCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGCCTGAAGATTT  
TGCAACTTATTACTGTCAACAGTTTAATAGTTTCCCGCTCACTTTCGGCGGAGGGACCA  
AGGTGGAGATCAAACGTACG

<The variable region of the light chain of HD8> (SEQ ID NO: 23)

[0147]

MDMRVPAQLLGLLLLWLPGARCAIQLTQSPSSLSASVGDRVTTTCRASQGISSALAWYQQ  
KPGKAPKLLIYDASSLESGVPSRFSGSGSGTDFLTISLQPEDFATYYCQQFNSFPLTFGGG  
TKVEIKRTV

[0148] The translation initiation site in the heavy chain DNA is the ATG codon starting from the 41st adenine (A) from the 5'-terminus of the sequence as shown in SEQ ID NO: 20, and the boundary between the variable region and the constant region of the antibody is located between the 496th adenine (A) and the 497th guanine (G) from the 5'-terminus. In the amino acid sequence, the variable region of the heavy chain is the portion between the N-terminus and the 152nd serine (S) residue in the sequence as shown in SEQ ID NO: 21, and the portion comprising the 153rd alanine (A) and thereafter is the constant region. The N-terminus of the purified heavy chain protein was analyzed. This demonstrated that the signal sequence of the heavy chain was the portion between the N-terminus and the 20th serine (S) of the sequence as shown in SEQ ID NO: 21, and the N-terminus of the matured body was the 21st glutamine (Q) of the sequence as shown in SEQ ID NO: 21. Accordingly, the matured portion in the amino acid sequence as shown in SEQ ID NO: 21 is the portion between the 21st glutamic acid and the 152nd serine.

[0149] The translation initiation site in the light chain DNA is the ATG codon starting from the 34th A from the 5'-

terminus of the sequence as shown in SEQ ID NO: 22, and the variable region is the portion between the 5'-terminus and the 420th adenine (A). In the amino acid sequence, the variable region is the portion between the N-terminus and the 129th lysine (K) of the sequence as shown in SEQ ID NO: 23. The N-terminus of the purified light chain protein was analyzed. This demonstrated that the signal sequence of the light chain is the portion between the N-terminus and the 22nd cysteine (C) of the sequence as shown in SEQ ID NO: 23, and the N-terminus of the matured body was the 23rd adenine (A) of the sequence as shown in SEQ ID NO: 23. Accordingly, the matured portion in the amino acid sequence as shown in SEQ ID NO: 23 is the portion between the 23rd adenine and the 129th lysine.

**Table 2 Nucleotide sequences of synthetic DNA**

Nucleotide sequences of synthetic DNA				
No	Primer	Sequence (5' to 3')	Length	SEQ ID NO.
1	hh-6	GGT CCG GGA GAT CAT GAG GGT GTC CTT	27	5
2	hh-3	GTG CAC GCC GCT GGT CAG GGC GCC TG	26	6
3	hh-4	GGT GCC AGG GGG AAG ACC GAT GG	23	7
4	tnHD4Sal	ata tgt cga cCC AGC CCT GGG ATT TTC AGG TGT TTT C	37	8
5	tnHD8Sal	ata tgt cga cTGG CTG ACC AGG GCA GTC ACC AGA G	35	9
6	tnCHNhe	gat ggg ccc ttg gtg cta gct gag gag acg g	31	10
7	hk-2	GTT GAA GCT CTT TGT GAC GGG CGA GC	26	11
8	hk-6	T GGC GGG AAG ATG AAG ACA GAT GGT G	26	12
9	tnHD4Bgl	ata tag atc tGC TGC TCA GTT AGG ACC CAG AGG GAA CC	38	13
10	tnHD8Bgl	ata tag atc tGG GAG TCA GAC CCA CTC AGG ACA CAG C	37	14
11	tnCkBsi	aag aca gat ggt gca gcc acc gta cgt ttg at	32	15

#### Example 13: Preparation of subclass recombinant vectors

**[0150]** The variable regions of the HD4 and HD8 antibodies were incorporated into the following variety of vectors. The N5KG1-Val Lark vector was used for the IgG1 type, the N5KG4 Lark vector was used for the IgG4 type (both vectors were manufactured by IDEC Pharmaceuticals, modified vectors of N5KG1 were disclosed in USP 6,001,358, N5KG1-Val Lark was modified while sustaining IgG1 as with N5KG1, and N5KG4 Lark was rearranged to the IgG4 type), a vector in which the constant region of the heavy chain of the N5KG1-Val Lark was rearranged into the IgG2 type (N5KG2) was used for the IgG2 type, and the variable regions of HD4 and HD8 were incorporated into the vector in the same manner as in Example 12. Separately, a vector in which the sequence CCC encoding proline (P) 331 according to the EU numbering system in the constant region of the heavy chain that was incorporated into the IgG1 or IgG2 vector was varied to TCC encoding serine (S) (Mi-Hua Tao et al., 1993, J. Exp. Med.) was prepared (hereafter referred to as N5KG1Ser and N5KG2Ser in that order), and the variable regions of HD4 and HD8 were incorporated into the vector in the same manner as in Example 12. In Fig. 3 and Table 3 below, for example, the HD4 antibody having an IgG1 subclass is referred to as HD4IgG1 or HD4G1, and the antibody that expresses the gene in which the sequence CCC encoding proline (P) 331 according to the EU numbering system in the constant region of the heavy chain was varied to TCC encoding serine (S) is referred to as HD4IgG1Ser or HD4G1 Ser.

**Table 3**

Names of recombinant vectors and produced recombinant antibodies					
Recombinant antibody and its cytotoxic activity					
Antibody	Vector	Subclass	Recombinant antibody	ADCC activity	CDC activity
HD4	N5KG1-Val Lark	IgG1	HD4G1	+	+



Table 3 (continued)

Names of recombinant vectors and produced recombinant antibodies					
Recombinant antibody and its cytotoxic activity					
Antibody	Vector	Subclass	Recombinant antibody	ADCC activity	CDC activity
HD4	N5KG2Ser	IgG2Ser	HD4G2Ser	-	-
HD4	N5KG4 Lark	IgG4	HD4G4	-	-
HD8	N5KG1-Val Lark	IgG1	HD8G1	+	+
HD8	N5KG1Ser	IgG1Ser	HD8G1Ser	+	-
HD8	NSKG2	IgG2	HD8G2	-	+
HD8	N5KG2Ser	IgG2Ser	HD8G2Ser	-	-
HD8	NSKG4 Lark	IgG4	HD8G4	-	-

Example 14: Preparation of recombinant antibody

**[0151]** The recombinant antibody-expressing vectors constructed in Examples 12 and 13 were introduced into host cells to prepare a recombinant antibody-expressing cell. Examples of host cells for expression that can be used include dhfr-deficient CHO cells (ATCC CRL-9096), CHO-Ras (Katakura Y, et al., Cytotechnology, 31: 103-109, 1999), and HEK293T (ATCC CRL-11268).

**[0152]** Vectors were introduced into host cells by electroporation, lipofection, or the like. Electroporation was carried out linearizing about 2 µg of an antibody-expressing vector with a restriction enzyme, introducing the genes into 4 x 10<sup>6</sup> CHO cells using Bio-Rad electrophoretor at 350 V and 500 µF, and sowing them on a 96-well culture plate. Lipofection was carried out using LipofectAMINE Plus (Gibco BRL) in accordance with the instructions therefor. After introducing the vectors, an agent that corresponds to the selection marker for the expression vector was added, and the culture was continued. After the colonies were confirmed, the antibody-expressing cells were selected by the method described in Example 4. Antibodies were purified from the selected cells in accordance with Example 8.

Example 15: Examination of cytotoxic activity

**[0153]** Cytotoxic activities through an antibody were assayed. These activities comprised cytotoxic activity on the target cell in the presence of the NK cell or a cell with killer activity such as a neutrophil and an antibody, i.e., antibody-dependent cellular cytotoxicity (ADCC), and cytotoxic activity on the target cell in the presence of a complement and an antibody, i.e., complement-dependent cytotoxicity (CDC). The hybridoma-derived HD4 and HD8 antibodies prepared in Example 8 and recombinant antibodies derived from each of the CHO cells prepared in Example 14 were used. In this case, hlgG was used as a control.

**[0154]** In simple terms, radioactive chromium (Cr<sup>51</sup>) was incorporated into the cytoplasm of the target cell, and the amount of Cr<sup>51</sup> released in the culture solution due to the cell death was measured based on the γ dose.

**[0155]** More specifically, 10<sup>6</sup> Burkitt's lymphoma cells, Raji (ATCC CCL-86), as the target cells were suspended in 15 µL of fetal calf serum (FCS), 50 µL (37 MBq/mL) of Cr<sup>51</sup>-labeled sodium chromate (Perkin Elmer, hereafter referred to as "Cr<sup>51</sup>") was added, and culture was then conducted at 37°C for 1 hour. Subsequently, 10 mL of medium was added, centrifugation was carried out, and the medium was discarded. This procedure was repeated three times to remove Cr<sup>51</sup> that was not incorporated in the cell.

**[0156]** In the ADCC assay, 200,000 mononuclear cells, which were derived from the peripheral blood of a healthy person obtained by the method described in Example 6, relative to 2,000 Cr<sup>51</sup>-labeled target cells (total volume: 200 µL) were cultured in a round-bottom, 96-well plate (Falcon) together with antibodies at each concentration level at 37°C in the presence of 5% CO<sub>2</sub> for 4 hours.

**[0157]** In the CDC assay, human serum-derived complements (final concentration: 5%, SIGMA) relative to 2,000 Cr<sup>51</sup>-labeled target cells (total volume: 200 µL) were cultured in a round-bottom, 96-well plate together with antibodies at each concentration level at 37°C in the presence of 5% CO<sub>2</sub> for 2 hours.

**[0158]** In both of the ADCC and CDC assays, the plate was subjected to centrifugation after the culture in order to deposit cells. Thereafter, 50 µL of supernatant was transferred to a powder scintillator-containing 96-well plate (Luma-plate™-96, Packard) and dehydrated at 55°C for 1.5 hours. After confirming the dehydration, the plate was covered with a dedicated-purpose cover (TopSeal™-A, 96-well Microplates, Packard), and the γ dose was measured using a scintillation counter (TopCount, Packard).

[0159] The results are shown in Fig. 3 and Table 3. HD8IgG1Ser, HD8IgG2Ser, HD8IgG4, HD4G2Ser, and HD4G4 had no CDC activity, and HD8IgG1, HD8IgG2, and HD4G1 had CDC activity. Only HD8IgG1, HD8IgG1Ser, and HD4G1 had ADCC activity.

#### 5 Example 16: Effects of recombinant antibody on tumor-bearing mouse model

[0160] Similar models as used in Example 11 were used to examine the pharmacological function of the HD8 recombinant antibody using tumor-bearing mouse models.

[0161] At the outset, 5-week-old C.B-17/CR-SCID mice (CLEA Japan, Inc.) were purchased, anti-asialo GM1 antiserum (Wako Chemicals) was diluted, and 10  $\mu$ L each thereof was intravenously administered to each of the mice when they were 6 weeks old. On the next day, Burkitt's lymphoma cells Raji (ATCC CCL-86) were intravenously administered in amounts of  $5 \times 10^6$  cells per mouse. Three days after the Raji transplantation, each antibody was administered once in tail veins of mice in amounts of 0.1  $\mu$ g or 1  $\mu$ g per mouse. The number of surviving mice after the transplantation was observed.

[0162] The HD8G1Ser and HD8G2Ser prepared in Example 14 were administered once in amounts of 0.1 or 1  $\mu$ g per mouse. As an antibody negative control, a group of mice to which the hlgG antibody used as a negative control in Example 9 was administered at 1  $\mu$ g/head was provided. As a positive control, a group of mice to which HD8G1 was administered at 1  $\mu$ g/mouse was provided.

[0163] The results of the above experiment are shown in Fig. 4. All the samples died within 16 days after the Raji transplantation in the negative control, i.e., the hlgG-administered group. The number of surviving mice 20 days later is as follows. With the administration of 1  $\mu$ g per mouse, all 6 samples survived except for the hlgG group (Fig. 4A), and with the administration of 0.1  $\mu$ g per mouse, all 6 samples in the HD8G2Ser group and 2 samples in the HD8G1Ser group survived (Fig. 4B). The number of surviving mice 25 days later is as follows. With the administration of 1  $\mu$ g per mouse, 2 mice in the HD8G1Ser group, 1 mouse in the HD8G1 group, and 4 mice in the HD8G2Ser group survived (Fig. 4A), and with the administration of 0.1  $\mu$ g per mouse, 1 mouse only in the HD8G2Ser group survived (Fig. 4B).

[0164] These results demonstrated that the HD8 recombinant antibodies, HD8G1Ser and HD8G2Ser, also exhibited anti-tumor effects at very low dosages as with HD8. Although HD8G2Ser had neither of ADCC or CDC activity, it exhibited beneficial effects on animal models.

#### 30 Example 17: Epitope analyses of HD4 HD6, and HD8

[0165] The epitope of each antibody was analyzed by Western blotting in accordance with a conventional technique. In simple terms, a cellulose or PVDF membrane was blocked with Block Ace (Yukijirushi), etc., each antibody was allowed to react therewith at 1  $\mu$ g/mL as a primary antibody and at 0.5  $\mu$ g/mL as a secondary antibody using HRP conjugated anti rabbit IgG (DAKO), the HRP-labeled anti-human antibody (e.g., DAKO) was allowed to react therewith, and the chemiluminescence was detected using a chemiluminescent reagent (e.g., ECL Western blotting detection reagent, Amersham Bioscience) and a chemiluminescence detector (e.g., LAS-1000, Fuji Film).

(1) A membrane fraction was extracted from the HLA-DR-expressing lymphoma cell SKW 6.4 (ATCC TIB-215), and the HLA-DR protein was purified from the anti-HLA-DR antibody (K28N, the name of produced cell: mouse-mouse hybridoma K28, deposited internationally at the International Patent Organism Depositary of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) as of February 22, 1994 under the accession number of FERM BP-4577) using an affinity column. The obtained protein was boiled using 4-20% gradient gel (Daiichi Pure Chemicals Co., Ltd.) under nonreducing conditions (95°C, 5 minutes), electrophoresis was carried out at a constant current of 25 mA for 1.5 hours per gel, and the resultant was transferred to a PVDF membrane at a constant current of 150 mA for 1 hour per gel. Subsequently, Western blot analysis was carried out in accordance with a conventional technique, and as a result, all of HD4, HD6, and HD8 were found to recognize the HLA-DR  $\beta$  chain located at approximately 30 Kda.

(2) Subsequently, regarding 199 amino acids in the extracellular region of the HLA-DR  $\beta$  chain (DRB1\*15011) (199 amino acids from amino acids 29 to 227 in the amino acid sequence as shown in SEQ ID NO: 147 and the nucleotide sequence encoding the amino acid sequence as shown in SEQ ID NO: 147 is shown in SEQ ID NO: 146), 94 types of peptides in total (SEQ ID NOs: 52 to 145) were spot synthesized from the C terminus on the cellulose membrane by shifting two amino acids in the 13-mer peptides, and the N terminus was acetylated (JERINI, Germany). The subsequent reaction was carried out based on conventional Western blot analysis (see, for example, Reineke, U. et al., (2001), "Epitope mapping with synthetic peptides prepared by SPOT synthesis." Antibody Engineering (Springer Lab Manual) Eds.: Kontermann/Dubel, 433-459). In the analysis, LumimagerTM (Boehringer-Mannheim) was used to represent the color intensity in each spot by numerical values.

The results are shown in Fig. 5. HD4 (Fig. 5A) exhibited potent reactivity with amino acids 61 to 73 and low

reactivity with amino acids 17 to 29, 63 to 75, and 65 to 77, and maximally bound to a peptide having the amino acid sequence as shown in SEQ ID NO: 83. HD6 (Fig. 5B) exhibited potent reactivity with amino acids 61 to 73 and low reactivity with amino acids 57 to 69 and 59 to 71, and maximally bound to a peptide having the amino acid sequence as shown in SEQ ID NO: 83. HD8 (Fig. 5C) exhibited very potent reactivity with amino acids 61 to 73, 63 to 75, and 65 to 77 and potentially bound to at least one peptide having the amino acid sequences as shown in SEQ ID Nos: 82, 83, and 84. In contrast, based on the conformation of HLA-DR (see, for example, Dessen A et al., *Immunity* (1997), 7, 473-481), amino acids 61 to 73 form the  $\alpha$  helix structure at the site where antigen-presenting peptides are retained.

(3) The  $\beta$ -chain polymorphisms of the 13 amino acids (amino acids 61 to 73) that exhibit maximal reactivity with all of HD4, HD6, and HD8 were taken into consideration, and 16 types of peptides including substantially almost all of the currently known about 350 types of polymorphisms (see the IMGT/HLA database of EMBL-EBI, etc.) were prepared. Further, 12 peptides (SEQ ID Nos: 40 to 51) in which each amino acid has been substituted with alanine (guanidine if it is originally alanine) were subjected to Western blotting under similar conditions. In the analysis, LAS2000 and ImageGauge analyzing software (Fuji Photo Film) were used to represent the color intensity in each spot by numerical values. Fig. 6 shows 28 types of peptide sequences and their reactivity with HD4, HD6, and HD8. The sequence to which each antibody positively reacted was represented by + to ++++ depending on its intensity, and represented by - in case of negative reactivity.

**[0166]** Positive or negative reaction was judged in accordance with the following criteria. Less than 5 % of the background: -

5% to less than 10% of the background: +/-  
 10% to less than 20% of the background: +  
 20% to less than 30% of the background: ++  
 30% to less than 50% of the background: +++  
 50% or more of the background: ++++

**[0167]** HD8 positively reacted with all sequences except for those as shown in SEQ ID Nos: 48 to 51 in which amino acids 65, 66, 69, and 72 had been substituted with alanine. The amino acids 65, 66, 69, and 72 are conserved in substantially almost all of the discovered HLA-DR  $\beta$  chains (see the IMGT/HLA database of EMBL-EBI, etc.) Since they cover most of the HLA-DR  $\beta$  chain sequences including major sequences among those as shown in SEQ ID Nos: 24 to 39, HD8 is very highly likely to be a Pan-HLA-DR antibody that binds to substantially almost all the HLA-DR  $\beta$  chains.

**[0168]** Furthermore, reactivity of HD8 was examined using the HLA-DR positive cell strain in the same manner as in Example 10. As a result, it was found to react with ARH77 (ATCC CRL-1621), Daudi (ATCC CCL-213), HS-Sultan (ATCC CRL-1484), IM-9 (ATCC CCL-159), MC/CAR (ATCC CRL-8083), Raji (ATCC CCL-86), Ramos (ATCC CRL-1596), RL (ATCC CRL-2261), SKW6.4 (ATCC TIB-215), and the L-cell (ATCC CCL-1) in which DRB1\*15011/DRA\*0101 were forcibly expressed and 5 specimens of human peripheral blood mononuclear cells derived from a healthy Japanese person. It has not yet been discovered in a cell in which nonreactive HLA-DR is expressed. Further, it reacts with 15 out of 15 crab-eating monkey specimens and 1 out of 1 chimpanzee specimen. In contrast, HD8 did not react with RPMI 8226 (ATCC CCL-155), which is a B-cell strain in which HLA-DR is not expressed.

#### Industrial Applicability

**[0169]** The present invention provides a preventive or therapeutic agent for diseases caused by HLA-DR-expressing cells, and more particularly, a molecule that is useful as a therapeutic agent for malignant tumors for patients having substantially almost all HLA-DR polymorphisms.

**[0170]** Further, the present invention provides an immunosuppressive agent for suppressing immunological activity associated with HLA-DR, and more particularly, a molecule that is useful as a therapeutic agent for rheumatisms.

**[0171]** All publications cited herein are incorporated herein by reference in their entirety. A person skilled in the art would easily understand that various modifications and changes are possible within the technical idea and the scope of the invention as described in the attached claims. The present invention is intended to include such modifications and changes.

SEQUENCE LISTING

5

<110> KIRIN BEER KABUSHIKI KAISHA

10

<120> Anti HLA-DR antibody

15

<130> PH-1646-PCT

20

<140>

<141>

25

<150> JP2001/317054

<151> 2001-10-15

30

<160> 147

35

<170> PatentIn Ver. 2.1

40

<210> 1

<211> 38

<212> DNA

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence:primer

50

<400> 1

55

ccggaattcc caccatggcc ataagtggag tccctgtg

38

5       <210> 2  
       <211> 38  
       <212> DNA  
 10       <213> Artificial Sequence  
  
 15       <220>  
       <223> Description of Artificial Sequence:primer  
  
 20       <400> 2  
       aaagcggccg ctcattacag aggccccctg cgttctgc               38  
  
 25  
  
       <210> 3  
 30       <211> 33  
       <212> DNA  
       <213> Artificial Sequence  
  
 35       <220>  
       <223> Description of Artificial Sequence:primer  
  
 40       <400> 3  
       ccggaattcc tggtcctgtc ctgttctcca gca               33  
 45  
  
 50       <210> 4  
       <211> 38  
       <212> DNA  
 55       <213> Artificial Sequence

5           <220>  
             <223> Description of Artificial Sequence:primer

10           <400> 4  
             aaagcggccg ctcatcagct caggaatcct gtiggctg                               38

15

            <210> 5  
 20           <211> 27  
             <212> DNA  
             <213> Artificial Sequence

25

            <220>  
 30           <223> Description of Artificial Sequence:primer

            <400> 5  
 35           ggtcgggag atcatgaggg tgcctt   27

40

            <210> 6  
             <211> 26  
 45           <212> DNA  
             <213> Artificial Sequence

50

            <220>  
             <223> Description of Artificial Sequence:primer

55

            <400> 6

gtgcacgccg ctggtcaggg cgcctg

26

5

<210> 7

10

<211> 23

<212> DNA

<213> Artificial Sequence

15

<220>

20

<223> Description of Artificial Sequence:primer

<400> 7

25

ggtgccaggg ggaagaccga tgg

23

30

<210> 8

<211> 37

35

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence:primer

45

<400> 8

atatgtcgac ccagccctgg gattttcagg tgttttc

37

50

55

<210> 9

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 9

atatgtcgac tggctgacca gggcagtcac cagag

35

<210> 10

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 10

gatgggccct tggctgctagc tgaggagacg g

31

<210> 11

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer



*5*           <400> 11  
gttgaagctc ttttgacgg gcgagc                         *26*

10

15           <210> 12  
            <211> 26  
            <212> DNA  
            <213> Artificial Sequence

20           <220>  
            <223> Description of Artificial Sequence:primer

25

<400> 12  
 30 tggcggaag atgaagacag atggtg 26

30

35	$\langle 210 \rangle$	13
	$\langle 211 \rangle$	38
	$\langle 212 \rangle$	DNA
40	$\langle 213 \rangle$	Artificial Sequence

35

&lt;211&gt; 38

&lt;212&gt; DNA

40

### ⟨213⟩ Artificial Sequence

45  $\langle 220 \rangle$

45

<223> Description of Artificial Sequence:primer

50  $\langle 400 \rangle$  13

50

atatagatct gctgctcagt taggaccag agggaacc 38

55

<210> 14

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 14

atatagatct gggagtcaga ccactcagg acacagc

37

<210> 15

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 15

aagacagatg gtgcagccac cgtacgtttg at

32

<210> 16

<211> 510

<212> DNA

<213> Homo sapiens

&lt;400&gt; 16

5 gtcgaccag cccctgggatt ttcaggtgtt ttcaggtgtt ttcatttggg gatcaggact 60  
 gaacagagag aactcaccat ggagtttggg ctgagctggc tttttcttgt ggctatttta 120  
 aaaggtgtcc agtgtgaggt gcaactgttg gagtctgggg gaggccttgg acagcctggg 180  
 10 gggctccctga gactctcctg tgcagcctct ggattcacct ttagcagcta tgccatgacc 240  
 tgggtccgcc aggcctccagg gaaggggctg gatlgggtct caggtattag tgggtgggtg 300  
 gatagcacat actacgcaga ctccgtgaag ggccgggtca ccatctccag agacaattcc 360  
 15 aagaacacgc tgtatctgca aatgaacagc ctgagagccg aggacacggc cgtatattac 420  
 tgtgcgagag atcatggttc ggggagttat tatccctact ggtttgacta ctggggccag 480  
 20 ggaaccctgg tcaccgtctc ctcagctagc 510

25  
 <210> 17

&lt;211&gt; 143

30  
 <212> PRT

&lt;213&gt; Homo sapiens

35  
 <400> 17

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala Ile Leu Lys Gly

40 1 5 10 15

Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln

45 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe

50 35 40 45

Ser Ser Tyr Ala Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu

55 50 55 60

5 Glu Trp Val Ser Gly Ile Ser Gly Gly Gly Asp Ser Thr Tyr Tyr Ala  
65 70 75 80

10 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn  
85 90 95

15 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val  
100 105 110

20 Tyr Tyr Cys Ala Arg Asp His Gly Ser Gly Ser Tyr Tyr Pro Tyr Trp  
115 120 125

25 Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala  
130 135 140

35 <210> 18

<211> 424

<212> DNA

40 <213> Homo sapiens

45 <400> 18

agatctgctg ctcagttagg acccagaggg aaccatggaa accccagcgc agcttctctt 60  
cctcctgcta ctcaggctcc cagataccac cggagaactt ggttgacgc agtctccagg 120  
50 caccctgtct ttgtctccag gggaaagagc caccctctcc tgcagggccca gtcagagtgt 180  
tagcagccgc tacttagcct ggtaccagca gaaacctggc caggctccca ggctcctcat 240  
ctatggtgca tccagcaggg ccactggcat cccagacagg ttcagtggca gtgggtcttg 300  
55 gacagacttc actctcacca tcagcagact ggagcctgaa gatcttgcag tgtattactg 360

tcagcagtat ggtagctcac cgctcacttt cggcggaggg accaaggtagg agatcaaacg 420

tacg 424

<210> 19

<211> 130

<212> PRT

<213> Homo sapiens

<400> 19

Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro

1 5 10 15

Asp Thr Thr Gly Glu Leu Val Leu Thr Gln Ser Pro Gly Thr Leu Ser

20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser

35 40 45

Val Ser Ser Arg Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala

50 55 60

Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro

65 70 75 80

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile

85 90 95

Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr

100

105

110

5

Gly Ser Ser Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys

115

120

125

10

Arg Thr

15

130

20

&lt;210&gt; 20

&lt;211&gt; 502

&lt;212&gt; DNA

25

&lt;213&gt; Homo sapiens

30

&lt;400&gt; 20

gtcgactggc tgaccagggc agtcaccaga gctccagaea atgtctgtct ccttccatcat 60  
 cttccctgccc gtgctgggcc tcccatgggg tgctctgtca caggttcagc tgcagcactc 120  
 aggtccagga ctggtgaagc cctcgagac cctctcactc acctgtgcca tctccgggga 180  
 cagtgtctct agcaacagt ctctcttgaa ctggatcagg cagtccecat cgaggggcct 240  
 tgagtggctg ggaaggacat actacaggic caagtggtat aatgattatg cagtatctgt 300  
 gaaaagtcga atagtcatca acccagacac atccaagaac cagttctccc tgcagctgaa 360  
 ctctgtgact cccgaggaca cggctgtgta ttactgtgcg agagaaaatt tctatggttc 420  
 ggagacttgt cataagaagt attactgcta cggatatggac gtctggggcc aagggaccac 480  
 ggtcaccgtc tcttcagcta gc 502

50

&lt;210&gt; 21

&lt;211&gt; 154

55

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

5

&lt;400&gt; 21

Met Ser Val Ser Phe Leu Ile Phe Leu Pro Val Leu Gly Leu Pro Trp  
 10 1 5 10 15

Gly Val Leu Ser Gln Val Gln Leu Gln His Ser Gly Pro Gly Leu Val  
 15 20 25 30

Lys Pro Ser Gln Thr Leu Ser Leu Thr Cys Ala Ile Ser Gly Asp Ser  
 20 35 40 45

Val Ser Ser Asn Ser Ala Ser Trp Asn Trp Ile Arg Gln Ser Pro Ser  
 25 50 55 60

Arg Gly Leu Glu Trp Leu Gly Arg Thr Tyr Tyr Arg Ser Lys Trp Tyr  
 30 65 70 75 80

Asn Asp Tyr Ala Val Ser Val Lys Ser Arg Ile Val Ile Asn Pro Asp  
 35 85 90 95

Thr Ser Lys Asn Gln Phe Ser Leu Gln Leu Asn Ser Val Thr Pro Glu  
 40 100 105 110

Asp Thr Ala Val Tyr Tyr Cys Ala Arg Glu Asn Phe Tyr Gly Ser Glu  
 45 115 120 125

Thr Cys His Lys Lys Tyr Tyr Cys Tyr Gly Met Asp Val Trp Gly Gln  
 50 130 135 140

Gly Thr Thr Val Thr Val Ser Ser Ala Ser

145

150

<210> 22

<211> 426

<212> DNA

<213> Homo sapiens

<400> 22

agatctggga gtcagaccca ctcaggacac agcatggaca tgagggtccc cgctcagctc 60  
 ctggggcctc tgcctgctctg gctcccaggt gccagatgtg ccatccagtt gaccagctct 120  
 ccatectccc tgtctgcac tctaggagac agatcacca tcacttgccg ggcaagtcag 180  
 ggcattagca gtgcctttagc ctggatcag cagaaaccag ggaaagctcc taaactcctg 240  
 atctatgatg cctccagttt ggaaagtggg gtcccatcaa ggctcagcgg cagtggatct 300  
 gggacagatt tcactctcac catcagcagc ctgcagcctg aagattttgc aacttattac 360  
 tgtcaacagt ttaatagttt cccgctcact ttcggcggag ggaccaaggt ggagatcaaa 420  
 cgtacg 426

<210> 23

<211> 132

<212> PRT

<213> Homo sapiens

<400> 23

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp

1

5

10

15



5 Leu Pro Gly Ala Arg Cys Ala Ile Gln Leu Thr Gln Ser Pro Ser Ser  
20 25 30

10 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser  
35 40 45

15 Gln Gly Ile Ser Ser Ala Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys  
50 55 60

20 Ala Pro Lys Leu Leu Ile Tyr Asp Ala Ser Ser Leu Glu Ser Gly Val  
65 70 75 80

25 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr  
85 90 95

30 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln  
100 105 110

35 Phe Asn Ser Phe Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile  
115 120 125

40 Lys Arg Thr Val  
130

45  
50 <210> 24

<211> 13

55 <212> PRT

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:peptide

10

<400> 24

Trp Asn Ser Gln Lys Asp Phe Leu Glu Asp Arg Arg Ala

15

1

5

10

20

<210> 25

<211> 13

25

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:peptide

35

<400> 25

Trp Asn Ser Gln Lys Asp Phe Leu Glu Arg Arg Arg Ala

40

1

5

10

45

<210> 26

<211> 13

50

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:peptide

5

<400> 26

Trp Asn Ser Gln Lys Asp Phe Leu Glu Asp Glu Arg Ala

10

1

5

10

15

<210> 27

<211> 13

20

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:peptide

30

<400> 27

Trp Asn Ser Gln Lys Asp Phe Leu Glu Gln Ala Arg Ala

35

1

5

10

40

<210> 28

<211> 13

45

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:peptide

55

<400> 28

Trp Asn Ser Gln Lys Asp Ile Leu Glu Asp Glu Arg Ala

5           1                   5                   10

10           <210> 29

<211> 13

15           <212> PRT

<213> Artificial Sequence

20           <220>

<223> Description of Artificial Sequence:peptide

25           <400> 29

Trp Asn Ser Gln Lys Asp Ile Leu Glu Gln Lys Arg Gly

30           1                   5                   10

35           <210> 30

<211> 13

40           <212> PRT

<213> Artificial Sequence

45           <220>

<223> Description of Artificial Sequence:peptide

50           <400> 30

Trp Asn Ser Gln Lys Asp Ile Leu Glu Asp Arg Arg Ala

55           1                   5                   10

5       <210> 31  
       <211> 13  
       <212> PRT  
 10       <213> Artificial Sequence  
  
 15       <220>  
       <223> Description of Artificial Sequence:peptide  
  
 20       <400> 31  
       Trp Asn Ser Gln Lys Asp Ile Leu Glu Asp Arg Arg Gly  
       1                   5                   10  
 25  
  
 30       <210> 32  
       <211> 13  
       <212> PRT  
 35       <213> Artificial Sequence  
  
 40       <220>  
       <223> Description of Artificial Sequence:peptide  
  
 45       <400> 32  
       Trp Asn Ser Gln Lys Asp Ile Leu Glu Asp Lys Arg Ala  
       1                   5                   10  
 50  
  
 55       <210> 33  
       <211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 33

Trp Asn Ser Gln Lys Asp Ile Leu Glu Gln Ala Arg Ala  
1 5 10

<210> 34

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 34

Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Arg Arg Ala  
1 5 10

<210> 35

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence:peptide

<400> 35

10 Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Ala Arg Ala  
1 5 10

15

<210> 36

20 <211> 13

<212> PRT

25 <213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence:peptide

<400> 36

35 Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Lys Arg Gly  
1 5 10

40

<210> 37

45 <211> 13

<212> PRT

50 <213> Artificial Sequence

<220>

55 <223> Description of Artificial Sequence:peptide

<400> 37

Trp Asn Ser Gln Lys Asp Leu Leu Glu Asp Arg Arg Ala

1 5 10

<210> 38

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 38

Trp Asn Ser Gln Lys Asp Leu Leu Glu Arg Arg Arg Ala

1 5 10

<210> 39

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 39

Trp Asn Ser Gln Lys Asp Leu Leu Glu Asp Glu Arg Ala

1 5 10



5

&lt;210&gt; 40

&lt;211&gt; 13

10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

20

&lt;400&gt; 40

Trp Asn Ser Gln Lys Asp Ala Leu Glu Gln Arg Arg Ala

25

1

5

10

30

&lt;210&gt; 41

&lt;211&gt; 13

35

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

40

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

45

&lt;400&gt; 41

Trp Asn Ser Gln Lys Asp Leu Leu Glu Ala Arg Arg Ala

50

1

5

10

55

&lt;210&gt; 42

<211> 13

5 <212> PRT

<213> Artificial Sequence

10 <220>

<223> Description of Artificial Sequence:peptide

15 <400> 42

Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Ala Arg Ala

20 1 5 10

25 <210> 43

<211> 13

30 <212> PRT

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence:peptide

40 <400> 43

Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Arg Arg Gly

45 1 5 10

50 <210> 44

<211> 13

55 <212> PRT

<213> Artificial Sequence

5           <220>  
           <223> Description of Artificial Sequence:peptide  
  
 10           <400> 44  
           Ala Asn Ser Gln Lys Asp Leu Leu Glu Gln Arg Arg Ala  
           1                   5                   10

20           <210> 45  
           <211> 13  
           <212> PRT  
 25           <213> Artificial Sequence

30           <220>  
           <223> Description of Artificial Sequence:peptide  
  
 35           <400> 45  
           Trp Ala Ser Gln Lys Asp Leu Leu Glu Gln Arg Arg Ala  
           1                   5                   10

45           <210> 46  
           <211> 13  
           <212> PRT  
 50           <213> Artificial Sequence

55           <220>  
           <223> Description of Artificial Sequence:peptide

5 <400> 46

Trp Asn Ala Gln Lys Asp Leu Leu Glu Gln Arg Arg Ala

1 5 10

15 <210> 47

<211> 13

<212> PRT

20 <213> Artificial Sequence

25 <220>

<223> Description of Artificial Sequence:peptide

30 <400> 47

Trp Asn Ser Ala Lys Asp Leu Leu Glu Gln Arg Arg Ala

1 5 10

40 <210> 48

<211> 13

<212> PRT

45 <213> Artificial Sequence

50 <220>

<223> Description of Artificial Sequence:peptide

55 <400> 48

Trp Asn Ser Gln Ala Asp Leu Leu Glu Gln Arg Arg Ala

1 5 10

5

<210> 49

10

<211> 13

<212> PRT

15

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence:peptide

<400> 49

25

Trp Asn Ser Gln Lys Ala Leu Leu Glu Gln Arg Arg Ala

1 5 10

30

<210> 50

35

<211> 13

<212> PRT

40

<213> Artificial Sequence

<220>

45

<223> Description of Artificial Sequence:peptide

<400> 50

50

Trp Asn Ser Gln Lys Asp Leu Leu Ala Gln Arg Arg Ala

1 5 10

55

<210> 51

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 51

Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln Arg Ala Ala

1 5 10

<210> 52

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 52

Gly Asp Thr Arg Pro Arg Phe Leu Trp Gln Pro Lys Arg

1 5 10

<210> 53

<211> 13

<212> PRT

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:peptide

10

<400> 53

Thr Arg Pro Arg Phe Leu Trp Gln Pro Lys Arg Glu Cys

15

1

5

10

20

<210> 54

<211> 13

25

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:peptide

35

<400> 54

Pro Arg Phe Leu Trp Gln Pro Lys Arg Glu Cys His Phe

40

1

5

10

45

<210> 55

<211> 13

50

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:peptide

5

<400> 55

Phe Leu Trp Gln Pro Lys Arg Glu Cys His Phe Phe Asn

10

1 5 10

15

<210> 56

<211> 13

20

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:peptide

30

<400> 56

Trp Gln Pro Lys Arg Glu Cys His Phe Phe Asn Gly Thr

35

1 5 10

40

<210> 57

<211> 13

45

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:peptide

55

<400> 57



Pro Lys Arg Glu Cys His Phe Phe Asn Gly Thr Glu Arg

5           1                   5                   10

10           <210> 58

            <211> 13

15           <212> PRT

            <213> Artificial Sequence

20           <220>

            <223> Description of Artificial Sequence:peptide

25           <400> 58

Arg Glu Cys His Phe Phe Asn Gly Thr Glu Arg Val Arg

30           1                   5                   10

35           <210> 59

            <211> 13

40           <212> PRT

            <213> Artificial Sequence

45           <220>

            <223> Description of Artificial Sequence:peptide

50           <400> 59

Cys His Phe Phe Asn Gly Thr Glu Arg Val Arg Phe Leu

55           1                   5                   10

5           <210> 60  
            <211> 13  
            <212> PRT  
10           <213> Artificial Sequence  
  
15           <220>  
            <223> Description of Artificial Sequence:peptide  
  
20           <400> 60  
            Phe Phe Asn Gly Thr Glu Arg Val Arg Phe Leu Asp Arg  
25                1                           5                           10  
            .  
  
30           <210> 61  
            <211> 13  
            <212> PRT  
35           <213> Artificial Sequence  
  
40           <220>  
            <223> Description of Artificial Sequence:peptide  
  
45           <400> 61  
            Asn Gly Thr Glu Arg Val Arg Phe Leu Asp Arg Tyr Phe  
50                1                           5                           10  
            .  
  
55           <210> 62  
            <211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 62

Thr Glu Arg Val Arg Phe Leu Asp Arg Tyr Phe Tyr Asn

1 5 10

<210> 63

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 63

Arg Val Arg Phe Leu Asp Arg Tyr Phe Tyr Asn Gln Glu

1 5 10

<210> 64

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 64

Arg Phe Leu Asp Arg Tyr Phe Tyr Asn Gln Glu Glu Ser

1 5 10

<210> 65

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 65

Leu Asp Arg Tyr Phe Tyr Asn Gln Glu Glu Ser Val Arg

1 5 10

<210> 66

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 66

Arg Tyr Phe Tyr Asn Gln Glu Glu Ser Val Arg Phe Asp

1 5 10

<210> 67

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 67

Phe Tyr Asn Gln Glu Glu Ser Val Arg Phe Asp Ser Asp

1 5 10

<210> 68

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 68

Asn Gln Glu Glu Ser Val Arg Phe Asp Ser Asp Val Gly

1 5 10

5

&lt;210&gt; 69

&lt;211&gt; 13

10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

20

&lt;400&gt; 69

Glu Glu Ser Val Arg Phe Asp Ser Asp Val Gly Glu Phe

25

1

5

10

30

&lt;210&gt; 70

&lt;211&gt; 13

35

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

40

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

45

&lt;400&gt; 70

Ser Val Arg Phe Asp Ser Asp Val Gly Glu Phe Arg Ala

50

1

5

10

55

&lt;210&gt; 71

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 71

Arg Phe Asp Ser Asp Val Gly Glu Phe Arg Ala Val Thr

1 5 10

<210> 72

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 72

Asp Ser Asp Val Gly Glu Phe Arg Ala Val Thr Glu Leu

1 5 10

<210> 73

<211> 13

<212> PRT

<213> Artificial Sequence

5           <220>  
          <223> Description of Artificial Sequence:peptide

10           <400> 73  
          Asp Val Gly Glu Phe Arg Ala Val Thr Glu Leu Gly Arg  
15           1                   5                   10

20           <210> 74  
          <211> 13  
          <212> PRT  
25           <213> Artificial Sequence

30           <220>  
          <223> Description of Artificial Sequence:peptide

35           <400> 74  
          Gly Glu Phe Arg Ala Val Thr Glu Leu Gly Arg Pro Asp  
40           1                   5                   10

45           <210> 75  
          <211> 13  
          <212> PRT  
50           <213> Artificial Sequence

55           <220>  
          <223> Description of Artificial Sequence:peptide



<400> 75

Phe Arg Ala Val Thr Glu Leu Gly Arg Pro Asp Ala Glu

1 5 10

<210> 76

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 76

Ala Val Thr Glu Leu Gly Arg Pro Asp Ala Glu Tyr Trp

1 5 10

<210> 77

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 77

Thr Glu Leu Gly Arg Pro Asp Ala Glu Tyr Trp Asn Ser

1 5 10

5

<210> 78

10

<211> 13

<212> PRT

15

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence:peptide

<400> 78

25

Leu Gly Arg Pro Asp Ala Glu Tyr Trp Asn Ser Gln Lys

1 5 10

30

<210> 79

35

<211> 13

<212> PRT

<213> Artificial Sequence

40

<220>

45

<223> Description of Artificial Sequence:peptide

<400> 79

50

Arg Pro Asp Ala Glu Tyr Trp Asn Ser Gln Lys Asp Ile

1 5 10

55

<210> 80

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 80

Asp Ala Glu Tyr Trp Asn Ser Gln Lys Asp Ile Leu Glu

1

5

10

<210> 81

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 81

Glu Tyr Trp Asn Ser Gln Lys Asp Ile Leu Glu Gln Ala

1

5

10

<210> 82

<211> 13

<212> PRT

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:peptide

10

<400> 82

Trp Asn Ser Gln Lys Asp Ile Leu Glu Gln Ala Arg Ala

15

1 5 10

20

<210> 83

<211> 13

25

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:peptide

35

<400> 83

Ser Gln Lys Asp Ile Leu Glu Gln Ala Arg Ala Ala Val

40

1 5 10

45

<210> 84

<211> 13

50

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:peptide

5

<400> 84

Lys Asp Ile Leu Glu Gln Ala Arg Ala Ala Val Asp Thr

10

1

5

10

15

<210> 85

<211> 13

20

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:peptide

30

<400> 85

Ile Leu Glu Gln Ala Arg Ala Ala Val Asp Thr Tyr Cys

35

1

5

10

40

<210> 86

<211> 13

45

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:peptide

55

<400> 86

Glu Gln Ala Arg Ala Ala Val Asp Thr Tyr Cys Arg His

5           1                   5                   10

10           <210> 87

            <211> 13

15           <212> PRT

            <213> Artificial Sequence

20           <220>

            <223> Description of Artificial Sequence:peptide

25           <400> 87

Ala Arg Ala Ala Val Asp Thr Tyr Cys Arg His Asn Tyr

30           1                   5                   10

35           <210> 88

            <211> 13

40           <212> PRT

            <213> Artificial Sequence

45           <220>

            <223> Description of Artificial Sequence:peptide

50           <400> 88

Ala Ala Val Asp Thr Tyr Cys Arg His Asn Tyr Gly Val

55           1                   5                   10

5           <210> 89  
             <211> 13  
             <212> PRT  
 10           <213> Artificial Sequence  
  
 15           <220>  
             <223> Description of Artificial Sequence:peptide  
  
 20           <400> 89  
             Val Asp Thr Tyr Cys Arg His Asn Tyr Gly Val Val Glu  
               1                   5                   10  
 25  
  
 30           <210> 90  
             <211> 13  
             <212> PRT  
 35           <213> Artificial Sequence  
  
 40           <220>  
             <223> Description of Artificial Sequence:peptide  
  
 45           <400> 90  
             Thr Tyr Cys Arg His Asn Tyr Gly Val Val Glu Ser Phe  
               1                   5                   10  
 50  
  
 55           <210> 91  
             <211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 91

Cys Arg His Asn Tyr Gly Val Val Glu Ser Phe Thr Val

1 5 10

<210> 92

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 92

His Asn Tyr Gly Val Val Glu Ser Phe Thr Val Gln Arg

1 5 10

<210> 93

<211> 13

<212> PRT

<213> Artificial Sequence



<220>

5 <223> Description of Artificial Sequence:peptide

<400> 93

10 Tyr Gly Val Val Glu Ser Phe Thr Val Gln Arg Arg Val  
1 5 10

15

<210> 94

20 <211> 13

<212> PRT

25 <213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence:peptide

<400> 94

35 Val Val Glu Ser Phe Thr Val Gln Arg Arg Val Gln Pro  
1 5 10

40

<210> 95

45 <211> 13

<212> PRT

50 <213> Artificial Sequence

<220>

55 <223> Description of Artificial Sequence:peptide

<400> 95

Glu Ser Phe Thr Val Gln Arg Arg Val Gln Pro Lys Val

1 5 10

<210> 96

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 96

Phe Thr Val Gln Arg Arg Val Gln Pro Lys Val Thr Val

1 5 10

<210> 97

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 97

Val Gln Arg Arg Val Gln Pro Lys Val Thr Val Tyr Pro

1 5 10

5

&lt;210&gt; 98

&lt;211&gt; 13

10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

20

&lt;400&gt; 98

Arg Arg Val Gln Pro Lys Val Thr Val Tyr Pro Ser Lys

25

1

5

10

30

&lt;210&gt; 99

&lt;211&gt; 13

35

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

40

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

45

&lt;400&gt; 99

Val Gln Pro Lys Val Thr Val Tyr Pro Ser Lys Thr Gln

50

1

5

10

55

&lt;210&gt; 100

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 100

Pro Lys Val Thr Val Tyr Pro Ser Lys Thr Gln Pro Leu

1 5 10

<210> 101

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 101

Val Thr Val Tyr Pro Ser Lys Thr Gln Pro Leu Gln His

1 5 10

<210> 102

<211> 13

<212> PRT

<213> Artificial Sequence

5           <220>  
          <223> Description of Artificial Sequence:peptide

10           <400> 102  
          Val Tyr Pro Ser Lys Thr Gln Pro Leu Gln His His Asn  
15           1                   5                   10

20           <210> 103  
          <211> 13  
          <212> PRT  
25           <213> Artificial Sequence

30           <220>  
          <223> Description of Artificial Sequence:peptide

35           <400> 103  
          Pro Ser Lys Thr Gln Pro Leu Gln His His Asn Leu Leu  
40           1                   5                   10

45           <210> 104  
          <211> 13  
          <212> PRT  
50           <213> Artificial Sequence

55           <220>  
          <223> Description of Artificial Sequence:peptide

5 <400> 104  
Lys Thr Gln Pro Leu Gln His His Asn Leu Leu Val Cys  
1 5 10  
10

15 <210> 105  
<211> 13  
<212> PRT  
20 <213> Artificial Sequence

25 <220>  
<223> Description of Artificial Sequence:peptide

30 <400> 105  
Gln Pro Leu Gln His His Asn Leu Leu Val Cys Ser Val  
1 5 10  
35

40 <210> 106  
<211> 13  
<212> PRT  
45 <213> Artificial Sequence

50 <220>  
<223> Description of Artificial Sequence:peptide

55 <400> 106  
Leu Gln His His Asn Leu Leu Val Cys Ser Val Ser Gly

1 5 10

5

<210> 107

10

<211> 13

<212> PRT

15

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence:peptide

<400> 107

25

His His Asn Leu Leu Val Cys Ser Val Ser Gly Phe Tyr

1 5 10

30

<210> 108

35

<211> 13

<212> PRT

<213> Artificial Sequence

40

<220>

45

<223> Description of Artificial Sequence:peptide

<400> 108

50

Asn Leu Leu Val Cys Ser Val Ser Gly Phe Tyr Pro Gly

1 5 10

55

<210> 109

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 109

Leu Val Cys Ser Val Ser Gly Phe Tyr Pro Gly Ser Ile

1 5 10

<210> 110

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 110

Cys Ser Val Ser Gly Phe Tyr Pro Gly Ser Ile Glu Val

1 5 10

<210> 111

<211> 13

<212> PRT



<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:peptide

10

<400> 111

Val Ser Gly Phe Tyr Pro Gly Ser Ile Glu Val Arg Trp

15

1 5 10

20

<210> 112

<211> 13

25

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:peptide

35

<400> 112

Gly Phe Tyr Pro Gly Ser Ile Glu Val Arg Trp Phe Leu

40

1 5 10

45

<210> 113

<211> 13

50

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:peptide

5

<400> 113

Tyr Pro Gly Ser Ile Glu Val Arg Trp Phe Leu Asn Gly

10

1 5 10

15

<210> 114

<211> 13

20

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:peptide

30

<400> 114

Gly Ser Ile Glu Val Arg Trp Phe Leu Asn Gly Gln Glu

35

1 5 10

40

<210> 115

<211> 13

45

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:peptide

55

<400> 115

Ile Glu Val Arg Trp Phe Leu Asn Gly Gln Glu Glu Lys

5           1                   5                           10

10           <210> 116

            <211> 13

15           <212> PRT

            <213> Artificial Sequence

20           <220>

            <223> Description of Artificial Sequence:peptide

25           <400> 116

Val Arg Trp Phe Leu Asn Gly Gln Glu Glu Lys Ala Gly

30           1                   5                           10

35           <210> 117

            <211> 13

40           <212> PRT

            <213> Artificial Sequence

45           <220>

            <223> Description of Artificial Sequence:peptide

50           <400> 117

Trp Phe Leu Asn Gly Gln Glu Glu Lys Ala Gly Met Val

55           1                   5                           10

5       <210> 118  
      <211> 13  
      <212> PRT  
10       <213> Artificial Sequence  
  
15       <220>  
      <223> Description of Artificial Sequence:peptide  
  
20       <400> 118  
      Leu Asn Gly Gln Glu Glu Lys Ala Gly Met Val Ser Thr  
      1                   5                   10  
25  
  
30       <210> 119  
      <211> 13  
      <212> PRT  
35       <213> Artificial Sequence  
  
40       <220>  
      <223> Description of Artificial Sequence:peptide  
  
45       <400> 119  
      Gly Gln Glu Glu Lys Ala Gly Met Val Ser Thr Gly Leu  
      1                   5                   10  
50  
  
55       <210> 120  
      <211> 13

<212> PRT

5 <213> Artificial Sequence

<220>

10 <223> Description of Artificial Sequence:peptide

<400> 120

15 Glu Glu Lys Ala Gly Met Val Ser Thr Gly Leu Ile Gln  
1 5 10

20

<210> 121

25 <211> 13

<212> PRT

30 <213> Artificial Sequence

<220>

35 <223> Description of Artificial Sequence:peptide

<400> 121

40 Lys Ala Gly Met Val Ser Thr Gly Leu Ile Gln Asn Gly  
1 5 10

45

<210> 122

50 <211> 13

<212> PRT

55 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 122

Gly Met Val Ser Thr Gly Leu Ile Gln Asn Gly Asp Trp

1

5

10

<210> 123

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 123

Val Ser Thr Gly Leu Ile Gln Asn Gly Asp Trp Thr Phe

1

5

10

<210> 124

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 124

Thr Gly Leu Ile Gln Asn Gly Asp Trp Thr Phe Gln Thr

1 5 10

<210> 125

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 125

Leu Ile Gln Asn Gly Asp Trp Thr Phe Gln Thr Leu Val

1 5 10

<210> 126

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 126

Gln Asn Gly Asp Trp Thr Phe Gln Thr Leu Val Met Leu

1 5 10

5

&lt;210&gt; 127

&lt;211&gt; 13

10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

20

&lt;400&gt; 127

Gly Asp Trp Thr Phe Gln Thr Leu Val Met Leu Glu Thr

25

1

5

10

30

&lt;210&gt; 128

&lt;211&gt; 13

35

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

40

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:peptide

45

&lt;400&gt; 128

Trp Thr Phe Gln Thr Leu Val Met Leu Glu Thr Val Pro

50

1

5

10

55

&lt;210&gt; 129



<211> 13

5 <212> PRT

<213> Artificial Sequence

10 <220>

<223> Description of Artificial Sequence:peptide

15 <400> 129

Phe Gln Thr Leu Val Met Leu Glu Thr Val Pro Arg Ser

20 1 5 10

25 <210> 130

<211> 13

30 <212> PRT

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence:peptide

40 <400> 130

Thr Leu Val Met Leu Glu Thr Val Pro Arg Ser Gly Glu

45 1 5 10

50 <210> 131

<211> 13

55 <212> PRT

<213> Artificial Sequence

5           <220>  
           <223> Description of Artificial Sequence:peptide  
  
 10          <400> 131  
           Val Met Leu Glu Thr Val Pro Arg Ser Gly Glu Val Tyr  
           1                   5                   10  
 15

20          <210> 132  
           <211> 13  
           <212> PRT  
 25          <213> Artificial Sequence

30          <220>  
           <223> Description of Artificial Sequence:peptide  
  
 35          <400> 132  
           Leu Glu Thr Val Pro Arg Ser Gly Glu Val Tyr Thr Cys  
           1                   5                   10  
 40

45          <210> 133  
           <211> 13  
           <212> PRT  
 50          <213> Artificial Sequence

55          <220>  
           <223> Description of Artificial Sequence:peptide

5 <400> 133

Thr Val Pro Arg Ser Gly Glu Val Tyr Thr Cys Gln Val

1 5 10

15 <210> 134

<211> 13

<212> PRT

20 <213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence:peptide

30 <400> 134

Pro Arg Ser Gly Glu Val Tyr Thr Cys Gln Val Glu His

1 5 10

40 <210> 135

<211> 13

<212> PRT

45 <213> Artificial Sequence

<220>

50 <223> Description of Artificial Sequence:peptide

55 <400> 135

Ser Gly Glu Val Tyr Thr Cys Gln Val Glu His Pro Ser

1 5 10

5

<210> 136

10

<211> 13

<212> PRT

15

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence:peptide

<400> 136

25

Glu Val Tyr Thr Cys Gln Val Glu His Pro Ser Val Thr

1 5 10

30

<210> 137

35

<211> 13

<212> PRT

<213> Artificial Sequence

40

<220>

45

<223> Description of Artificial Sequence:peptide

<400> 137

50

Tyr Thr Cys Gln Val Glu His Pro Ser Val Thr Ser Pro

1 5 10

55

<210> 138

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 138

Cys Gln Val Glu His Pro Ser Val Thr Ser Pro Leu Thr

1

5

10

<210> 139

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 139

Val Glu His Pro Ser Val Thr Ser Pro Leu Thr Val Glu

1

5

10

<210> 140

<211> 13

<212> PRT

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:peptide

10

<400> 140

15

His Pro Ser Val Thr Ser Pro Leu Thr Val Glu Trp Arg

1

5

10

20

<210> 141

<211> 13

25

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:peptide

35

<400> 141

Ser Val Thr Ser Pro Leu Thr Val Glu Trp Arg Ala Arg

40

1

5

10

45

<210> 142

<211> 13

50

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence:peptide

5

<400> 142

Thr Ser Pro Leu Thr Val Glu Trp Arg Ala Arg Ser Glu

10

1

5

10

15

<210> 143

<211> 13

20

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:peptide

30

<400> 143

Pro Leu Thr Val Glu Trp Arg Ala Arg Ser Glu Ser Ala

35

1

5

10

40

<210> 144

<211> 13

45

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:peptide

55

<400> 144

Thr Val Glu Trp Arg Ala Arg Ser Glu Ser Ala Gln Ser

5           1                   5                   10

10       <210> 145

<211> 13

15       <212> PRT

<213> Artificial Sequence

20       <220>

<223> Description of Artificial Sequence:peptide

25       <400> 145

Val Glu Trp Arg Ala Arg Ser Glu Ser Ala Gln Ser Lys

30           1                   5                   10

35       <210> 146

<211> 801

40       <212> DNA

<213> Homo sapiens

45       <400> 146

atggtgtgtc tgaagctccc tggaggctcc tgcattgacag cgctgacagt gacactgatg 60  
gtgctgagct cccactggc ttgtctggg gacacccgac cacgtttcct gtggcagcct 120  
50 aagagggagt gicatttcct caatgggacg gagcgggtgc ggttcctgga cagatacttc 180  
tataaccagg aggagtcctt gcgcttcgac agcgacgtgg gggagttccg ggcggtgacg 240  
gagctggggc ggcctgacgc tgagtactgg aacagccaga aggacatcct ggagcaggcg 300  
55 cgggccgcgg tggacacctt ctgcagacac aactacgggg ttgtggagag cttcacagtg 360



cagcggcgag tccaacctaa ggtgactgta tatecttcaa agaccagcc cctgcagcac 420  
 5 cacaacctcc tggctcgtc tctgagtgg tctatccag gcagcattga agtcaggtgg 480  
 ttcctgaacg gccaggaaga gaaggctggg atgggtgtcca caggcctgat ccagaatgga 540  
 gactggacct tccagaccct ggtgatgctg gaaacagttc ctggaagtgg agaggtttac 600  
 10 acctgccaag tggagcacc aagcgtgaca agccctctca cagtgggaatg gagagcacgg 660  
 tctgaatctg cacagagcaa gatgctgagt ggagtcgggg gctttgtgct gggcctgctc 720  
 15 ttccttgggg ccgggctggt catctacttc aggaatcaga aaggacactc tggacttcag 780  
 ccaacaggat tctgagctg a 801

20 <210> 147

<211> 266

25 <212> PRT

<213> Homo sapiens

30 <400> 147

Met Val Cys Leu Lys Leu Pro Gly Gly Ser Cys Met Thr Ala Leu Thr

35 1 5 10 15

Val Thr Leu Met Val Leu Ser Ser Pro Leu Ala Leu Ser Gly Asp Thr

40 20 25 30

45 Arg Pro Arg Phe Leu Trp Gln Pro Lys Arg Glu Cys His Phe Phe Asn

35 40 45

50 Gly Thr Glu Arg Val Arg Phe Leu Asp Arg Tyr Phe Tyr Asn Gln Glu

50 55 60

55 Glu Ser Val Arg Phe Asp Ser Asp Val Gly Glu Phe Arg Ala Val Thr

	65		70		75		80
5							
	Glu	Leu	Gly	Arg	Pro	Asp	Ile
				85		90	95
10							
	Leu	Glu	Gln	Ala	Arg	Ala	Tyr
				100		105	110
15							
	Gly	Val	Val	Glu	Ser	Phe	Val
				115		120	125
20							
	Thr	Val	Tyr	Pro	Ser	Lys	Thr
				130		135	140
25							
	Val	Cys	Ser	Val	Ser	Gly	Phe
				145		150	155
30							
	Phe	Leu	Asn	Gly	Gln	Glu	Glu
				165		170	175
35							
	Ile	Gln	Asn	Gly	Asp	Trp	Thr
				180		185	190
40							
	Val	Pro	Arg	Ser	Gly	Glu	Val
				195		200	205
45							
	Val	Thr	Ser	Pro	Leu	Thr	Val
				210		215	220
50							
	Val	Thr	Ser	Pro	Leu	Thr	Val
				210		215	220
55							

	Gln	Ser	Lys	Met	Leu	Ser	Gly	Val	Gly	Gly	Phe	Val	Leu	Gly	Leu	Leu
5	225					230					235					240

	Phe	Leu	Gly	Ala	Gly	Leu	Phe	Ile	Tyr	Phe	Arg	Asn	Gln	Lys	Gly	His
10					245					250					255	

	Ser	Gly	Leu	Gln	Pro	Thr	Gly	Phe	Leu	Ser
15				260					265	

20

25

30

35

40

45

50

55

**Claims**

1. An antibody that binds to HLA-DR or a functional fragment thereof, which is produced from the hybridoma HD8 (accession number FERM BP-7773).
2. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region of an antibody produced from the hybridoma HD8 (accession number: FERM BP-7773).
3. The antibody or a functional fragment thereof according to claim 2, wherein the antibody subclass is IgG.
4. The antibody or a functional fragment thereof according to claim 3, wherein IgG is IgG1.
5. The antibody or a functional fragment thereof according to claim 4, wherein an amino acid sequence in the constant region of the heavy chain is modified.
6. The antibody or a functional fragment thereof according to claim 5, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.
7. The antibody or a functional fragment thereof according to claim 3, wherein IgG is IgG2.
8. The antibody or a functional fragment thereof according to claim 7, wherein an amino acid sequence in the constant region of the heavy chain is modified.
9. The antibody or a functional fragment thereof according to claim 8, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.
10. The antibody or a functional fragment thereof according to claim 3, wherein IgG is IgG3.
11. The antibody or a functional fragment thereof according to claim 10, wherein an amino acid sequence in the constant region of the heavy chain is modified.
12. The antibody or a functional fragment thereof according to claim 3, wherein IgG is IgG4.
13. The antibody or a functional fragment thereof according to claim 12, wherein an amino acid sequence in the constant region of the heavy chain is modified.
14. The hybridoma HD8 (accession number FERM BP-7773).
15. An antibody that binds to HLA-DR or a functional fragment thereof, which is produced from the hybridoma HD4 (accession number: FERM BP-7771).
16. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region of an antibody produced from the hybridoma HD4 (accession number: FERM BP-7771).
17. The antibody or a functional fragment thereof according to claim 16; wherein the antibody subclass is IgG.
18. The antibody or a functional fragment thereof according to claim 17, wherein IgG is IgG1.
19. The antibody or a functional fragment thereof according to claim 18, wherein an amino acid sequence in the constant region of the heavy chain is modified.
20. The antibody or a functional fragment thereof according to claim 17, wherein IgG is IgG2.
21. The antibody or a functional fragment thereof according to claim 20, wherein an amino acid sequence in the constant region of the heavy chain is modified.

22. The antibody or a functional fragment thereof according to claim 21, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.
- 5 23. The antibody or a functional fragment thereof according to claim 17, wherein IgG is IgG3.
24. The antibody or a functional fragment thereof according to claim 23, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 10 25. The antibody or a functional fragment thereof according to claim 17, wherein IgG is IgG4.
26. The antibody or a functional fragment thereof according to claim 25, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 15 27. The hybridoma HD4 (accession number FERM BP-7771).
28. An antibody that binds to HLA-DR or a functional fragment thereof, which is produced from the hybridoma HD10 (accession number: FERM BP-7774).
- 20 29. The hybridoma HD10 (accession number FERM BP-7774).
30. An antibody that binds to HLA-DR or a functional fragment thereof, which is produced from the hybridoma HD6 (accession number: FERM BP-7772).
- 25 31. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region of an antibody produced from the hybridoma HD6 (accession number: FERM BP-7772).
32. The antibody or a functional fragment thereof according to claim 31, wherein the antibody subclass is IgG.
- 30 33. The antibody or a functional fragment thereof according to claim 32, wherein IgG is IgG1.
34. The antibody or a functional fragment thereof according to claim 33, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 35 35. The antibody or a functional fragment thereof according to claim 32, wherein IgG is IgG2.
36. The antibody or a functional fragment thereof according to claim 35, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 40 37. The antibody or a functional fragment thereof according to claim 32, wherein IgG is IgG3.
38. The antibody or a functional fragment thereof according to claim 37, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 45 39. The antibody or a functional fragment thereof according to claim 32, wherein IgG is IgG4.
40. The antibody or a functional fragment thereof according to claim 39, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 50 41. The hybridoma HD6 (accession number FERM BP-7772).
42. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 21 and 23.
- 55 43. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region comprising the mature variable regions of peptides encoded by the nucleotide sequences as shown in SEQ ID NOs: 20 and 22.

44. The antibody or a functional fragment thereof according to claim 42, wherein the antibody subclass is IgG.
45. The antibody or a functional fragment thereof according to claim 44, wherein IgG is IgG1.
- 5 46. The antibody or a functional fragment thereof according to claim 45, wherein an amino acid sequence in the constant region of the heavy chain is modified.
47. The antibody or a functional fragment thereof according to claim 46, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.
- 10 48. The antibody HD8G1Ser or a functional fragment thereof, which is the antibody HD8 having an IgG1 subclass and amino acid 331 according to the EU numbering system being substituted with Ser.
- 15 49. The antibody or a functional fragment thereof according to claim 44, wherein IgG is IgG2.
50. The antibody or a functional fragment thereof according to claim 49, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 20 51. The antibody or a functional fragment thereof according to claim 50, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.
- 25 52. The antibody HD8G2Ser or a functional fragment thereof, which is the antibody HD8 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser.
53. The antibody or a functional fragment thereof according to claim 44, wherein IgG is IgG3.
54. The antibody or a functional fragment thereof according to claim 53, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 30 55. The antibody or a functional fragment thereof according to claim 44, wherein IgG is IgG4.
56. The antibody or a functional fragment thereof according to claim 55, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 35 57. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 17 and 19.
- 40 58. An antibody that binds to HLA-DR or a functional fragment thereof, which comprises a variable region comprising the mature variable regions of peptides encoded by the nucleotide sequences as shown in SEQ ID NOs: 16 and 18.
59. The antibody or a functional fragment thereof according to claim 57, wherein the antibody subclass is IgG.
- 45 60. The antibody or a functional fragment thereof according to claim 59, wherein IgG is IgG1.
61. The antibody or a functional fragment thereof according to claim 60, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 50 62. The antibody or a functional fragment thereof according to claim 59, wherein IgG is IgG2.
63. The antibody or a functional fragment thereof according to claim 62, wherein an amino acid sequence in the constant region of the heavy chain is modified.
- 55 64. The antibody or a functional fragment thereof according to claim 63, wherein the modification of the amino acid sequence in the constant region of the heavy chain is the substitution of amino acid 331 according to the EU numbering system with Ser.

65. The antibody HD4G2Ser or a functional fragment thereof, which is the antibody HD4 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser.
66. The antibody or a functional fragment thereof according to claim 59, wherein IgG is IgG3.
67. The antibody or a functional fragment thereof according to claim 66, wherein an amino acid sequence in the constant region of the heavy chain is modified.
68. The antibody or a functional fragment thereof according to claim 59, wherein IgG is IgG4.
69. The antibody or a functional fragment thereof according to claim 68, wherein an amino acid sequence in the constant region of the heavy chain is modified.
70. An antibody that binds to HLA-DR or a functional fragment thereof, which maximally binds to the peptide as shown in SEQ ID NO: 82 selected from among peptides prepared by shifting 2 amino acids of the amino acids in the extracellular region of the HLA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides, binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus.
71. An antibody that binds to HLA-DR or a functional fragment thereof, which potently binds to all three peptides as shown in SEQ ID NOs: 82, 83, and 84 selected from among peptides prepared by shifting 2 amino acids of the amino acids in the extracellular region of the HLA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides, binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus.
72. An antibody that binds to HLA-DR or a functional fragment thereof, which significantly binds to all the peptides as shown in SEQ ID NOs: 24 to 39 and all the peptides as shown in SEQ ID NOs: 40 to 43 prepared by shifting 2 amino acids of the amino acids in the extracellular region of the HLA-DR  $\beta$  chain (DRB1\*15011) to prepare 13-mer peptides, binding the resulting peptides to a cellulose membrane through the C-terminus, and acetylating the N-terminus.
73. An antibody that binds to HLA-DR or a functional fragment thereof, which has the following properties (a) and (b):
- (a) when a 6-week-old SCID mouse is inoculated intravenously with 10  $\mu$ l of the anti-asialo GM1 antiserum, on the next day, inoculated intravenously with  $5 \times 10^6$  of Burkitt's lymphoma cells Raji (ATCC CCL-86), and 5 days thereafter, inoculated once with 5  $\mu$ g/kg, based on body weight, of the antibody of the present invention, the survival ratio of the mouse 90days after the inoculation is higher than that 90 days after the inoculation with the same amount of the human anti-HSA antibody; and
- (b) the immunosuppressive activity is lower than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC HB-55) at the same concentration, wherein the immunosuppressive activity is assayed as follows: 50  $\mu$ l of antibody adjusted at 8 mg/mL using the 10% FCS-containing RPMI 1640 medium is mixed with 50  $\mu$ L of mature dendritic cell suspension derived from a first human donor adjusted at  $2 \times 10^5$  cells/mL using the 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate, the mixture is allowed to stand at 4°C for 30 minutes, the resultant is mixed with 100  $\mu$ L of T-cell suspension (purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using the 10% FCS-containing RPMI 1640 medium derived from a second human donor having a histocompatible antigen different from that of the first human donor, the mixture is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 5 days, <sup>3</sup>H thymidine is added thereto at 1.0  $\mu$ Ci/well, the resultant is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 16 to 20 hours, the <sup>3</sup>H thymidine incorporated in the cell is recovered and then measured using a scintillator, and the incorporation of the <sup>3</sup>H thymidine into the cell is used as an indicator to assay the immunosuppressive activity.
74. The antibody or a functional fragment thereof according to claim 73, wherein the antibody is a monoclonal antibody.
75. The antibody or a functional fragment thereof according to claim 73, wherein the antibody is a human antibody.
76. The antibody or a functional fragment thereof according to claim 73, which is produced from a mouse-mouse hybridoma.
77. An antibody that binds to HLA-DR or a functional fragment thereof, which has the immunosuppressive activity equivalent to or higher than that achieved when using the mouse anti-HLA-DR monoclonal antibody L243 (ATCC

HB-55) at the same concentration, wherein the immunosuppressive activity is assayed as follows: 50 µl of antibody adjusted at 8 mg/mL using the 10% FCS-containing RPMI 1640 medium is mixed with 50 µL of mature dendritic cell suspension derived from a first human donor adjusted at  $2 \times 10^5$  cells/mL using the 10% FCS-containing RPMI 1640 medium in wells of a 96-well plate, the mixture is allowed to stand at 4°C for 30 minutes, the resultant is mixed with 100 µL of T-cell suspension (purity: 99% or higher) adjusted at  $1 \times 10^6$  cells/mL using the 10% FCS-containing RPMI 1640 medium derived from a second human donor having a histocompatible antigen different from that of the first human donor, the mixture is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 5 days, <sup>3</sup>H thymidine is added thereto at 1.0 µCi/well, the resultant is cultured at 37°C in the presence of 5% CO<sub>2</sub> for 16 to 20 hours, the <sup>3</sup>H thymidine incorporated in the cell is recovered and then measured using a scintillator, and the incorporation of the <sup>3</sup>H thymidine into the cell is used as an indicator to assay the immunosuppressive activity.

78. The antibody or a functional fragment thereof according to claim 77, wherein the antibody is a monoclonal antibody.

79. The antibody or a functional fragment thereof according to claim 77, wherein the antibody is a human antibody.

80. The antibody or a functional fragment thereof according to claim 77, which is produced from a mouse-mouse hybridoma.

81. A nucleic acid encoding an antibody comprising a variable region of an antibody produced from a hybridoma or a functional fragment thereof, wherein said nucleic acid is possessed by a hybridoma selected from the group consisting of the hybridoma HD8 (accession number FERM BP-7773), the hybridoma HD10 (accession number FERM BP-7774), the hybridoma HD4 (accession number FERM BP-7771), and the hybridoma HD6 (accession number FERM BP-7772).

82. A nucleic acid encoding the antibody or a functional fragment thereof according to claim 81, which comprises a variable region having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 17 and 19.

83. A nucleic acid encoding the antibody or a functional fragment thereof according to claim 81, which comprises a variable region having an amino acid sequence of the mature variable regions of the amino acid sequences as shown in SEQ ID NOs: 21 and 23.

84. A nucleic acid encoding the antibody or a functional fragment thereof, wherein the antibody is selected from the group consisting of the antibody HD8G1Ser, which is the antibody HD8 having an IgG1 subclass and amino acid 331 according to the EU numbering system being substituted with Ser, the antibody HD8G2Ser, which is the antibody HD8 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser, and the antibody HD4G2Ser, which is the antibody HD4 having an IgG2 subclass and amino acid 331 according to the EU numbering system being substituted with Ser.

85. A protein encoded by the nucleic acid according to any one of claims 81 to 84.

86. An expression vector having the nucleic acid according to any one of claims 81 to 84.

87. A host having the expression vector according to claim 86.

88. The host according to claim 87, which is selected from the group consisting of *E. coli*, yeast cell, insect cell, mammalian animal cell, plant cell, and mammalian animals.

89. A process for producing the anti-HLA-DR monoclonal antibody, wherein a gene encoding the anti-HLA-DR monoclonal antibody is isolated from a hybridoma selected from the group consisting of the hybridoma HD8 (accession number FERM BP-7773), the hybridoma HD10 (accession number FERM BP-7774), the hybridoma HD4 (accession number FERM BP-7771), and the hybridoma HD6 (accession number FERM BP-7772), an expression vector having said gene is constructed, the expression vector is introduced into a host to express the monoclonal antibody, and the anti-HLA-DR monoclonal antibody is collected from the resulting host, a culture supernatant of the host, or a secretion product of the host.

90. A preventive, therapeutic, or diagnostic agent for tumors, which comprises, as an active ingredient, the antibody or a functional fragment thereof according to any one of claims 1 to 26, 28, and 42 to 76.



91. The preventive, therapeutic, or diagnostic agent for tumors according to claim 90, wherein the tumor is at least one member selected from the group consisting of leukemia (including chronic lymphatic leukemia and acute lymphatic leukemia), lymphoma (including non-Hodgkin's lymphoma, Hodgkin's lymphoma, T-cell lymphoma, B-cell lymphoma, Burkitt's lymphoma, malignant lymphoma, diffuse lymphoma, and follicular lymphoma), myeloma (including multiple myeloma), breast cancer, colon cancer, kidney cancer, gastric cancer, ovarian cancer, pancreatic cancer, cervical cancer, endometrial cancer, esophageal cancer, liver cancer, head and neck squamous cancer, skin cancer, urinary tract cancer, prostate cancer, choriocarcinoma, pharyngeal cancer, laryngeal cancer, pleural tumor, arrhenoblastoma, endometrial hyperplasia, endometriosis, embryoma, fibrosarcoma, Kaposi's sarcoma, angioma, cavernous angioma, hemangioblastoma, retinoblastoma, spongiocytoma, neurofibroma, oligodendroglioma, medulloblastoma, neuroblastoma, neuroglioma, rhabdomyoblastoma, glioblastoma, osteogenic sarcoma, leiomyosarcoma, thyroid sarcoma, and Wilms tumor.

92. An immunosuppressive agent, which comprises, as an active ingredient, the antibody or a functional fragment thereof according to any one of claims 30 to 40 and 77 to 80.

93. A preventive, therapeutic, or diagnostic agent for autoimmune diseases or allergies, which comprises, as an active ingredient, the antibody or a functional fragment thereof according to any one of claims 30 to 40 and 77 to 81.

Fig. 1




Fig. 2A

Dosage:  $1 \mu\text{g}/\text{head}$




Fig. 2B


Dosage: 0.1  $\mu$ g/head

Fig. 3A

ADCC




Fig. 3B

CDC




Fig. 3C




Fig. 3D

CDC






Fig. 4A




Fig. 4B








Fig. 5C

HD8




Fig. 6

SEQ ID NO	61	62	63	64	65	66	67	68	69	70	71	72	73	HD4	HD6	HD8
24	W	N	S	Q	K	D	F	L	E	D	R	R	A	++++	+++	+++
25	W	N	S	Q	K	D	F	L	E	R	R	R	A	++	++++	++++
26	W	N	S	Q	K	D	F	L	E	D	E	R	A	++++	-	+++
27	W	N	S	Q	K	D	F	L	E	Q	A	R	A	++++	++	++++
28	W	N	S	Q	K	D	I	L	E	D	E	R	A	+++	-	++++
29	W	N	S	Q	K	D	I	L	E	Q	K	R	G	+/-	+	++++
30	W	N	S	Q	K	D	I	L	E	D	R	R	A	+	+/-	++++
31	W	N	S	Q	K	D	I	L	E	D	R	R	G	+/-	-	++++
32	W	N	S	Q	K	D	I	L	E	D	K	R	A	+/-	-	++++
33	W	N	S	Q	K	D	I	L	E	Q	A	R	A	+++	+/-	++++
34	W	N	S	Q	K	D	L	L	E	Q	R	R	A	++	+++	++++
35	W	N	S	Q	K	D	L	L	E	Q	A	R	A	++++	-	++++
36	W	N	S	Q	K	D	L	L	E	Q	K	R	G	+	+	+++
37	W	N	S	Q	K	D	L	L	E	D	R	R	A	+++	+/-	++++
38	W	N	S	Q	K	D	L	L	E	R	R	R	A	++	++++	++++
39	W	N	S	Q	K	D	L	L	E	D	E	R	A	+++	-	+++
40	W	N	S	Q	K	D	A	L	E	Q	R	R	A	+	+++	++++
41	W	N	S	Q	K	D	L	L	E	A	R	R	A	++++	++++	++++
42	W	N	S	Q	K	D	L	L	E	Q	A	R	A	+++	++	++++
43	W	N	S	Q	K	D	L	L	E	Q	R	R	G	+	+++	+++
48	W	N	S	Q	A	D	L	L	E	Q	R	R	A	-	+/-	+/-
49	W	N	S	Q	K	A	L	L	E	Q	R	R	A	-	++	+/-
50	W	N	S	Q	K	D	L	L	A	Q	R	R	A	-	++	-
51	W	N	S	Q	K	D	L	L	E	Q	R	A	A	-	+/-	-
44	A	N	S	Q	K	D	L	L	E	Q	R	R	A	+/-	+++	++++
45	W	A	S	Q	K	D	L	L	E	Q	R	R	A	+	++	++++
46	W	N	A	Q	K	D	L	L	E	Q	R	R	A	+	++	++++
47	W	N	S	A	K	D	L	L	E	Q	R	R	A	+	++	++++

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/10665

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl <sup>7</sup> C07K16/28, C12N5/10, C12N15/13, C12N1/21, C12N1/19, C12P21/08, A61K39/395  According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl <sup>7</sup> C07K16/28, C12N5/10, C12N15/13, C12N1/21, C12N1/19, C12P21/08, A61K39/395  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) REGISTRY (STN), CA (STN), MEDLINE (STN), WPI (DIALOG), BIOSIS (DIALOG), GenBank/EMBL/DBJ/GeneSeq, SwissProt/PIR/GeneSeq		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KOSTELNY, S.A. et al., Humanization and characterization of the anti-HLA-DR antibody 1D10. Int. J. Cancer 2001. Aug, Vol.93, No.4, pages 556 to 565	1-93
Y	NIMAN, H.L. et al., Generation of protein-reactive antibodies by short peptides is an event of high frequency: implications for the structural basis of immune recognition. Proc. Natl. Acad. Sci. USA. 1983, Vol.80, No.16, pages 4949 to 4953	1-93
Y	ULRICH, R.G. et al., Immune recognition of human major histocompatibility antigens: localization by a comprehensive synthetic strategy of the continuous antigenic sites in the first domain of HLA-DR2 $\beta$ chain. Eur. J. Immunol. 1987, Vol.17, No.4, pages 497 to 502	1-93
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
Date of the actual completion of the international search 16 January, 2003 (16.01.03)		Date of mailing of the international search report 28 January, 2003 (28.01.03)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)